

Product datasheet for MR228340

Ube2w (NM_001271016) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ube2w (NM_001271016) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Ube2w
Synonyms: 6130401J04Rik
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >MR228340 representing NM_001271016
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGTTGGGCCCCCGCGCGTGACGCGAGCACGGCGGCTCCGCCCCCTGCGTCTCGGGCTCGCTGGCCTT
 GGGGGGATGGTTTCATCATGGCGTCAATGCAGAAACGACTACAAAAGAACTGTTGGCTTGCAGAAATGA
 CCCACCTCCTGGAATGACTTTAAATGAAAAGAGTGTTCAAGATTCAATCACGCAGTGGATCGTAGACATG
 GAAGGTGCACCAGGAACCTTATATGAAGGGGAAAAATTTCACTTTTGTAAATTTAGTAGTCGATACC
 CTTTGTACTCTCCTCAGAAAGACTGGTCCCCGGCGCTCTCAGTGCAGTCAGTCTGTCTCAGCATTATCAGC
 ATGCTTTCCAGCTGCAAAGAAAAGAGACGACCACAGATAATTCTTTTATGTGCGAACATGTAACAAGA
 ATCCAAAGAAAACAAAATGGTGGTATCATGGTGGATACAGTGCCCGTGGAGGCCAGAAGAGGGACTCTTA
 CCCACTGGAAGTGGAGTTTAAAGATAGTTGTGTGCTGTCATGTGAGTTCTATGGTCACC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR228340 representing NM_001271016
 Red=Cloning site Green=Tags(s)

MLGPRGVTRARRLRPLRLGPRWPWGDGFIASMQKRLQKELLALQNDPPPGMTLNEKSVQNSITQWIVDM
 EGAPGTLYEGEKFQLLFKFSRYPFDSPQKTGPRRSQCSQSVSALSACFPAAKKRDDHQIIPFMCEHVTR
 IQRKQNGGIMVDTPVEARRGTLTHWNWSLKIVCCHVSSMVT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI



Cloning Scheme:



ACCN: NM_001271016

ORF Size: 549 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_001271016.1](#), [NP_001257945.1](#)


RefSeq Size: 1551 bp

RefSeq ORF: 552 bp

Locus ID: 66799

Cytogenetics:	1 A3
MW:	21.2 kDa
Gene Summary:	<p>Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. Specifically monoubiquitinates the N-terminus of various substrates, including ATXN3, MAPT/TAU, POLR2H/RPB8 and STUB1/CHIP, by recognizing backbone atoms of disordered N-termini (PubMed:21855799, PubMed:21229326). Involved in degradation of misfolded chaperone substrates by mediating monoubiquitination of STUB1/CHIP, leading to recruitment of ATXN3 to monoubiquitinated STUB1/CHIP, and restriction of the length of ubiquitin chain attached to STUB1/CHIP substrates by ATXN3 (PubMed:21855799). After UV irradiation, but not after mitomycin-C (MMC) treatment, acts as a specific E2 ubiquitin-conjugating enzyme for the Fanconi anemia complex by associating with E3 ubiquitin-protein ligase FANCL and catalyzing monoubiquitination of FANCD2, a key step in the DNA damage pathway (PubMed:21229326). In vitro catalyzes 'Lys-11'-linked polyubiquitination. UBE2W-catalyzed ubiquitination occurs also in the presence of inactive RING/U-box type E3s, i.e. lacking the active site cysteine residues to form thioester bonds with ubiquitin, or even in the absence of E3, albeit at a slower rate (By similarity).[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR228340