

Product datasheet for **MR229128**

Wdr61 (NM_023191) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Wdr61 (NM_023191) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Wdr61
Synonyms:	2700038L12Rik; 2810418I05Rik; REC14
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR229128 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGC**C

ATGACCAACCACTACAGTATTCTCTTCAAGCAAGAGCAAGCCCATGATGATGCCATATGGTCAGTTGCCT
 GGGAGACAAACAAAAGGAAACATTGAAACAGTGGTCACAGGATCCCTGGATGACCTGGTGAAGGTCTG
 GAAATGGCGTGATGAGAGGCTGGAGCTCCAGTGGAGCCTGGAGGGACATCAGCTCGGGGTGGTGTCTGTG
 GACATCAGCCACACTCTCCCATTTGCTGCATCCAGCTCTCTAGACGCTCATATTCGCCTCTGGGACTTGG
 AAAATGGCAAACAGATGAAGTCTATAGATGCAGGACCAGTGGATGCCTGGACTTTGGCATTCTCTCTGA
 CTCCCAGTATCTGGCCACAGGAACTCACATGGGGAAGTGAACATTTTTGGTGTGGAAAGTGGAAAAAA
 GAATATTTCTTTGGACACTAGAGGAAAATTCATCCTTAGTATTGCATATAGTCCTGATGGGAAATACCTGG
 CCAGCGGAGCCATAGACGGAATCATCAATATTTTGGACATTGCAACTGGAAAGCTTTTGCATACGCTGGA
 AGGCCATGCGATGCCCATTCGCTCCTTGACCTTTCCCTGACTCCCAGCTCCTTGTCACGGCTTCAGAT
 GATGGCTACATCAAGATCTATGATGTACAACATGCCAATTTGGCTGGCACACTGAGTGGCCATGCGTCTT
 GGGTGTGAATGTTGCGTTCTGTCCTGATGACACTCACTTTGTCTCCAGTTCATCTGACAAAAGTGTGAA
 GGTTTGGGATGTTGGAACAAGGACCTGTATTCACACCTTCTTTGATCACCAGGATCAGGTTTGGGAGTA
 AAATAAATGAAATGGATCAAAAATTGTATCTGTTGGAGATGACCAGGAAATTCATGTCTATGACTGCC
 CAATT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)

Protein Sequence: >MR229128 protein sequence
 Red=Cloning site Green=Tags(s)


MTNQYSILFKQEQAHDDAIWSVAWETNKKENIETVVTGSLDDLKVKWKWRDERLELQWSLEGHQLGVVSV
 DISHTLPIAASSSLDAHRLWDLENGQMKSIDAGPVDATLAFSPDSQYLATGTHMGKVNIFGVESGKK
 EYSLDTRGKFILSIAYSPDGKYLASGAIDGIINIFDIATGKLLHTLEGHAMPIRSLTFSPDSQLLVTASD
 DGYIKIYDVQHANLAGTLSGHASWVLNVAFCPDDTHFVSSSSDKSVKVDVGTRTCIHTFFDHQDQVWGV
 KYNGNGSKIVSVGDDQEIHVYDCPI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_023191

ORF Size: 915 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_023191.3](#), [NP_075680.1](#)

RefSeq Size: 1225 bp

RefSeq ORF: 918 bp

Locus ID: 66317


UniProt ID: [Q9ERF3](#)

Cytogenetics: 9 A5.3

MW: 33.8 kDa

Gene Summary: Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Required for mono- and trimethylation on histone H3 'Lys-4' (H3K4me3), dimethylation on histone H3 'Lys-79' (H3K4me3). Required for Hox gene transcription. Component of the SKI complex which is thought to be involved in exosome-mediated RNA decay and associates with transcriptionally active genes in a manner dependent on PAF1C (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR229128