

Product datasheet for MR229341

Rassf1 (NM_001243748) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Rassf1 (NM_001243748) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Rassf1
Synonyms:	123F2; AA536941; AU044980; NORE2A; Rassf1A; Rassf1B; Rassf1C; RDA32; REH3P21
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>MR229341 representing NM_001243748 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGTCGGCGGAGCCAGAACTCATTGAACTACGCGAGCTGGCACCCTCCGGGCGCATCGGTCCAGGCCGTA
CCCGGCTGGAGCGTGCCAACGCTCTGCGCATCGCCCCGGGTACAACACGCAATCCGTACAGCAGCACGT
CCCGGCTCGTGGCCACCCTTTCCAGCCTGCAGGGCCCACCACGCACACGTGGTGGCAGCTCTGTGGAGAC
TTCATCTGGGGCGTCGTGCGCAAGGGCCTACAGTGCAGCGCACTGCAAGTTCACCTGCCATTACCGTTGTC
GCGCGCTCGTCTGCTTGGACTGCTGCGGGCCCCGAGACCTGGGCTGGGACTCTGCACTAGAGCGGGATAC
GAACGTGGATGAGGCTGTAGAGCGGGAGACACCCGATCTTTCTCAAGCTGAGACTGAGCAGAAAAATCAAG
GACTACAATGGCCAGATCAACAGCAACCTTTCATGAGCCTGAATAAGGATGGCTCCTACACAGGCTTCA
TCAAGGTTTCAGCTGAAACTAGTGCCTCTGTTTCAGTGCCTTCCAGCAAGAAACACCTTCTTACAGGA
CGCCCCGAGAGGTACGGGGCGGAGCACAGCCGTGAAGCGCCGACCTCTTTTACTTGCCTAAGGATGCT
ATTAAGCATTCGATGTTCTATCACGAACACGGGCACGTGAGGTCATTGAGGCCCTGCTTCGAAAATTCA
TGGTTCGATAGATCCTCGAAGTTTGCACCTTTGAGCGAAGTGAACGGCATGGCCAAAGTATACCTCCG
GAAGCTGTCGGATGACGAGCAGCCCTTGAAGCTGCGGCTTCTTGCAGGGCCAGTGAAAAAGCCCTGAGC
TTTGTCTGAAGGAAAATGACTCGGGAGAGGTGAACTGGGATGCCTTTCAGCATGCCTGAACTGCACAATT
TCTACGAATCCTGCAGCGGGAAGAAGAGGAACACCTTCGCCAGATCCTGCAGAAGTATTCTCGTTGTGC
CCAGAAGATCCAGGAGGCTCTGCACGCTGTCCTTTGGG

ACGGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR229341 representing NM_001243748
 Red=Cloning site Green=Tags(s)

MSAEPELIELRELAPSGRIGPGRTRLERANALRIAPGTTRNPSQQHVPGRGHRFQAGPTTHTWCDCD
 CGD FIWGVVRKGLQCAHCKFTCHYRCRALVCLDCCGPRDLGWDSALERDTNVDEAVERETPDLSQAETE
 QKIK DYNGQINSNLFMSLNKDGSYTGF IKVQLKLRPVSVPSSKKPPSLQDARRGTGRSTAVKRRTS
 FYLPKDA IKHLHVL SRTRAREVIEALLRKF MVVDDPRK FALFERTERHGQVYLRKLSDDDEQPLK
 LRLLAGPSEKALS FVLKENDSGEVNWDASFMPELHNFLRILQREEEHLRQILQKYSRCRQKI
 QEALHACPLG


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001243748

ORF Size: 1020 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001243748.1 , NP_001230677.1
RefSeq Size:	1787 bp
RefSeq ORF:	1023 bp
Locus ID:	56289
UniProt ID:	Q99MK9
Cytogenetics:	9 F1
MW:	39.2 kDa
Gene Summary:	Potential tumor suppressor. Required for death receptor-dependent apoptosis. Mediates activation of Mediates activation of STK3/MST2 and STK4/MST1 during Fas-induced apoptosis by preventing their dephosphorylation. When associated with MOAP1, promotes BAX conformational change and translocation to mitochondrial membranes in response to TNF and TNFSF10 stimulation. Isoform A interacts with CDC20, an activator of the anaphase-promoting complex, APC, resulting in the inhibition of APC activity and mitotic progression. Inhibits proliferation by negatively regulating cell cycle progression at the level of G1/S-phase transition by regulating accumulation of cyclin D1 protein. Isoform C has been shown not to perform these roles, no function has been identified for this isoform. Isoform A disrupts interactions among MDM2, DAXX and USP7, thus contributing to the efficient activation of TP53 by promoting MDM2 self-ubiquitination in cell-cycle checkpoint control in response to DNA damage (By similarity).[UniProtKB/Swiss-Prot Function]