

## **Product datasheet for MR229380**

## Fst (NM\_001301373) Mouse Tagged ORF Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Fst (NM\_001301373) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Fst

Synonyms: AL033346; D2Mgi5; FS

Mammalian Cell Neomycin

Selection:


Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 

Cloning sites used for ORF Shuttling:





		ECOR V							Flag. Tag							F	'me I	rse i	
(	TAE	CTG	GCA	GCA	AAT	GAT	ATC	CTG	GAT	TAC	AAG	GAT	GAC	GAC	GAT	AAG	GTT	TAA	ACGGCCGGCC
_	D	L	A	A	N	D	I	L	D	Y	K	D	D	D	D	K	v	Stop	

<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_001301373

ORF Size: 1032 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001301373.1</u>, <u>NP 001288302.1</u>

 RefSeq Size:
 2575 bp

 RefSeq ORF:
 1035 bp

 Locus ID:
 14313

 UniProt ID:
 P47931

 Cytogenetics:
 13 D2.2

MW: 38.3 kDa


Gene Summary: The protein encoded by this gene binds to and negatively regulates activin, as well as other

members of the transforming growth factor beta family, and acts to prevent uncontrolled cellular proliferation. This protein also contains a heparin-binding sequence. It is expressed in many of the tissues in which activin is synthesized and is thought to clear activin from the circulation by attachment to the cell surface. Alternative splicing results in multiple transcript variants that encode multiple protein isoforms, including FST315 and FST288, that differ at their C-terminus. Another isoform, FST303 is thought to be produced by proteolytic cleavage of FST315. These isoforms differ in their localization and in their ability to bind heparin. While FST315 is a circulating protein, FST288 is tissue-bound, and FST303 is gonad-specific. While deletion of all isoforms results in embryonic lethality, expression of just FST288 is sufficient for embryonic development, but the resultant mice have fertility defects. [provided by

RefSeq, Aug 2014]



## **Product images:**



Circular map for MR229380