

## Product datasheet for **MR230052**

### **Foxa2 (NM\_001291065) Mouse Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Foxa2 (NM_001291065) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Foxa2
Synonyms:	Hnf-3b; HNF3-beta; Hnf3b; HNF3beta; Tcf-3b; Tcf3b
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

**ORF Nucleotide Sequence:**

>MR230052 representing NM\_001291065  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGCACTCGGCTTCCAGTATGCTGGGAGCCGTGAAGATGGAAGGGCAGGACCCATCCGACTGGAGCAGCT  
 ACTACGGGAGCCCGAGGGCTACTCTCCGTGAGCAACATGAACGCCGGCCTGGGGATGAATGGCATGAA  
 CACATACATGAGCATGTCCGCGGCTGCCATGGGCGGCGGTTCCGGCAACATGAGCGCGGGTCCATGAAC  
 ATGTCATCCTATGTGGCGCTGGAATGAGCCCGTCGCTAGCTGGCATGTCCCGGGCGCCGGCGCCATGG  
 CGGGCATGAGCGGCTCAGCCGGGCGGCGGCGTGGCGGCATGGGACCTCACCTGAGTCCGAGTCTGAG  
 CCCGCTCGGGGACAGGCGGCGGGGCCATGGGTGGCCTTGCCCCCTACGCCAACATGAACCTGATGAGC  
 CCCATGTACGGGAGGCCGGCTGAGCCGCGCTCGGGACCCCAAGACATACCGACGCAGCTACACACACG  
 CCAAACCTCCCTACTCGTACATCTCGCTCATCACCATGGCCATCCAGCAGAGCCCCAACAGATGCTGAC  
 GCTGAGCGAGATCTATCAGTGGATCATGGACCTTCCCTTTCTACCGGAGAACCAGCAGCGCTGGCAG  
 AACTCCATCCGCCACTCTCTCTCTTCAACGACTGCTTTCTCAAGGTGCCCGCTCGCCAGACAAGCCTG  
 GCAAGGGCTCCTTCTGGACCCTGCACCCAGACTCGGGCAACATGTTGAGAACCGGCTGCTACCTGCGCCG  
 CCAGAAGCGCTTCAAGTGTGAGAAGCAACTGGCACTGAAGGAAGCCGCGGGTGGCGCCAGTAGCGGAGGC  
 AAGAAGACCGCTCCTGGGTCCCAGGCCTCTCAGGCTCAGCTCGGGGAGGCCGCGGGCTCGGCCTCCGAGA  
 CTCCGGCGGGCACCGAGTCCCCCATCCAGCGCTTCTCCGTGTCAGGAGCACAAGCGAGGTGGCCTAAG  
 CGAGCTAAAGGGAGCACCTGCCTCTGCGCTGAGTCTCCCGAGCCGGCGCCCTCGCTGGCAGCAGCAG  
 CAGGCTGCAGCCACCTGCTGGGCCACCTCACCACCCAGGCCTGCCACCAGAGGCCACCTGAAGCCCG  
 AGCACCATTACGCCTTCAACCACCCCTTCTCTATCAACAACCTCATGTCGTCCGAGCAGCAACATCACCA  
 CAGCCACCACCACCATCAGCCCCACAAAATGGACCTCAAGGCCTACGAACAGGTGATGCACTACCCAGGG  
 GGCTATGGTTCGCCATGCCAGGAGCTTGGCCATGGGCCAGTACGAACAAAGCGGGCTGGATGCTC  
 CGCCCTGGCTGCAGACACTTCTACTACCAAGGAGTGTACTCCAGGCCTATTATGAACCTATCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**

>MR230052 representing NM\_001291065  
 Red=Cloning site Green=Tags(s)


MHSASSMLGAVKMEGHEPSDWSSYYAEPEGYSSVSNMNAAGLGMNGMNTYMSMSAAAMGGGSGNMSAGSMN  
 MSSYVAGMSPSLAGMSPGAGAMAGMSGAGAGVAGMGPPLSPSLSPGGQAAGAMGGLAPYANMNSMS  
 PMYQAGLSRARDPKTYRRSYTHAKPPYSYISLITMAIQQSPNKMLTLSEIYQWIMDLFPFYRQNRQWQ  
 NSIRHLSFNDCFLKVPSPDKPGKGSFWTLHPDSGNMFENGCYLRQKRFKCEKQLALKEAAGAASSGG  
 KKTAPGSQASQQLGEAAGSASETPAGTESPHSSASPCQEHKRGGLSELKGAPASALSPPEPAPSPGQQQ  
 QAAAHLGPPHPGLPPEAHLKPEHHYAFNHPFSINNLMSSEQQHHHSHHHHQPHKMDLKAYEQVMHYPG  
 GYGSPMPGSLAMGPVTNKAGLDASPLAADTSYYQGVYSRPIMNSS

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:**

SgfI-MluI

**Cloning Scheme:**

**Plasmid Map:**


**ACCN:** NM\_001291065

**ORF Size:** 1395 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001291065.1](#), [NP\\_001277994.1](#)

**RefSeq Size:** 2070 bp

**RefSeq ORF:** 1398 bp

**Locus ID:** 15376

**UniProt ID:** [P35583](#)

**Cytogenetics:** 2 73.38 cM

**MW:** 49.5 kDa

**Gene Summary:** Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). In embryonic development is required for notochord formation. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; Foxa1 and Foxa2 seem to have at least in part redundant roles. FOXA1 and FOXA2 are essential for hepatic specification. FOXA1 and FOXA2 are required for morphogenesis and cell differentiation during formation of the lung. FOXA1 and FOXA2 are involved in bile duct formation; they positively regulate the binding glucocorticoid receptor/NR3C1 to the IL6 promoter. FOXA1 and FOXA2 regulate multiple phases of midbrain dopaminergic neuron development; they regulate expression of NEUROG2 at the beginning of mDA neurogenesis and of NR4A2 and EN1 in immature mDA neurons. Modulates the transcriptional activity of nuclear hormone receptors; inhibits AR-mediated transcription from the LCN5 promoter. Binds to fibrinogen beta promoter and is involved in IL6-induced fibrinogen beta transcriptional activation. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta-cells and glucose homeostasis. In pancreatic beta cells activates transcription of potassium channel subunits KCNJ11 and ABCC8. Involved in regulation of fat metabolism; activates transcriptional programs of lipid metabolism and ketogenesis at low insulin state. Involved in transcriptional regulation of MUC2 in the intestine.[UniProtKB/Swiss-Prot Function]