

Product datasheet for **MR230973**

Rps6ka1 (NM_001285506) Mouse Tagged ORF Clone

Product data:

| | |
|--------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | Rps6ka1 (NM_001285506) Mouse Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | Rps6ka1 |
| Synonyms: | Mapkapk-1a; p90-Rsk1; p90rsk; p90Rsk1; p90S6K; Rsk; Rsk-1; Rsk1; S6K-alpha-1 |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| Cell Selection: | Neomycin |



[View online »](#)

ORF Nucleotide Sequence:

>MR230973 representing NM_001285506
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGCAAACACCCGCCGATTTCCCTAGGGTTGAGAGAGACTCCGGCTTCTATCCCAGAAAGGATGAGGCCA
 TCCTCAAGGAGATCTCCATCACACACCAGTCAAGGCTGGCTCTGAGAAGGCCGATCCATCCCAGTTTGA
 GCTCCTCAAGTTCTGGGCCAAGGATCCTTTGGCAAAGTCTTCTGGTACGCAAGGTCACCCGGCCTGAC
 AGTGGGCACTTGTATGCCATGAAAGTATTAAGAAGGCCACGCTGAAAGTGCGTGACCGTGTTCGGACCA
 AGATGGAGAGAGACATCCTCGCTGACGTGAACCACCCGTTCTGGTGAAGCTACATTATGCCTCCAGAC
 CGAGGGCAAGCTCTATCTTATTCTGGACTTCTGCGTGGTGGAGACCTGTTACACACGGCTCTCAAAGGAG
 GTCATGTTACAGAGGAGGATGTGAAGTTTACCTGGCTGAGCTGGCACTGGGCCTGGACCCTGCACA
 GCTTGGGCATTATTTACAGAGACCTCAAGCCTGAGAATATCCTTTGGATGAGGAGGGCCACATCAAAC
 CACTGACTTTGGCCTGAGCAAGGAGGCCATAGACCATGAGAAGAAGGCTTACTCCTTCTGCGGGACAGTG
 GAGTACATGGCGCCCAGGTTGTCAACCGCCAGGGTCACACCCACAGTGCAGACTGGTGGTCTATGGGG
 TGCTGATGTTTGAGATGTGACGGGCTCCCTGCCCTTCCAGGGGAAGGACCGGAAGGAGACCATGACCTT
 GATTTTGAAGGCGAAGCTAGGCATGCCCGATTTCTGAGCACGGAAGCCCAGAGCCTCTGCGGGCCCTG
 TTCAAGAGGAATCCTGCCAATCGGCTTGGCTCGGGCCCTGATGGGGCAGAGGAAATTAAGAGACATATCT
 TCTACTCCACCATTGACTGGAATAAGCTTACCAGGCTGAGATCAAGCCCCCTTCAAGCCGGCTGTGGC
 CCAGCCGACGACACCTTCTACTTTGATACCGAGTTCACGTACGCACACCCAGGATTCGCCAGGCATC
 CCTCCCAGTGTGGTCCCATCAGCTGTTCCGTGGCTTACGCTTCGTGGCACTGGTCTGATGGAGGACG
 ACGGCAAGCCTCGGACCACTCAGGCCCCCTACACTCGTGGTACAGAACTCCACGGGAAGAATTTGGT
 TTTCAAGTACGGCTACGTAGTAAAGGAGACGATCGGCGTGGGCTCCTACTCTGTGTGTAAGCGTTGTGTC
 CACAAGGCCACCAACATGGAGTATGCTGTCAAGGTCATCGACAAGAGCAAAGAGATCCCTCAGAAGAGA
 TTGAGATTCTTCTGCGGTATGGCCAGCACCCCAACATCATCACCCTGAAAGATGTGTACGATGATGGTAA
 GCACGTGTACCTGGTACAGAGCTGATGAGGGGCGGCGAGCTGCTGGATAAGATCCTACGGCAGAAATTC
 TTCTCAGAGCGGGAGGCCAGCTTCTGCTGCACACGATCAGCAAGACTGTGGAATACTTGCACTCTCAAG
 GGGTTGTCCACAGAGATCTCAAACCCAGTAACATCCTCTATGTGGATGAGTCTGGAAACCCCGAGTGCCT
 GCGCATCTGCGACTTTGGCTTTGCCAAGCAGCTACGGGCCGAGAACGGACTCCTCATGACACCTTGCTAC
 ACAGCCAACTTTGTGGCACCTGAGGTGCTGAAGCGCCAGGGCTACGATGAAGGCTGTGACATATGGAGCC
 TGGGCATTCTGCTGTACACGATGCTGGCAGGATACACTCCATTTGCCAATGGGCCAGTGACACCCAGA
 GGAGATCCTCACCCGGATCGGCAGCGGGAAGTTACCCCTCAGTGGGGGAAACTGGAACACGGTTTCAGAG
 ACAGCCAAGGACTTGGTATCTAAGATGCTGCATGTGGACCCACCAGCGCCTCACAGCCAAGCAGGTCC
 TGCAGCACCCATGGATCACCCAGAAAGACAAGCTTCCCCAGAGCCAGTTGTCCCACCAAGACCTGCAGCT
 CGTGAAGGGAGCCATGGCGGCTACATACTCTGCTCTCAATAGCTCCAAACCCACCCCTCAGCTCAAGCCA
 ATTGAGTCGTCTATCCTGGCCAGCGGCGGGTGAAGGAGCTGCCATCCACCACCTG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR230973 representing NM_001285506
 Red=Cloning site Green=Tags(s)

MQTPADFP RVERDSGFYPRKDEAILKEISITHVVKAGSEKADPSQFELLKVLGQGSFGKVFVLRKVTRPD
 SGHLYAMKVLKKATLKVRDRVRTKMERDILADVNHPFVVKLHYAFQTEGKLYLILDFLRGGDLFTRL SKE
 VMFTEEDVKFYLAELALGLDHLHSLGIIYRDLKPENILLDEEGHIKLTDFGLSKEAIDHEKKAYSFCGTV
 EYMAPEVVNRQGHTHSADWWSYGVLMFEMLTGSLPFQGKDRKETMTLILKAKLGMPQFLSTEAQSLLRAL
 FKRNPANRLGSGPDGAEEIKRHIFYSTIDWNKLYRREIKPPFKPAVAQPDDTFYFDTEFTSRTPRDSPGI
 PPSAGAHQLFRGFSFVATGLMEDDGKPRTTQAPLHSHVQQLHGKNLVFSDGYVVKETIGVGSYSVCKRCV
 HKATNMEYAVKVIDKSKRDPSEEIEILLRYGQHPNIITLKD VYDDGKHVYLVTELMRGGELLDKILRQKF
 FSERESFVLHTISKTV EYLHSQGVVHRDLKPSNILYVDESGNPECLRICDFGFAKQLRAENGLLMTPCY
 TANFVAPEVLKRQGYDEGCDIWSL GILLYTMLAGYTPFANGPSDTP EEILTRIGSGKFTLSGGNWNVTSE
 TAKDLVSKMLHVDPHQRLTAKQVLQHPWITQKDKLPQSQLSHQDLQLVKGAMAATYSALNSSKPTPQLKP
 IESSILAQRVRKLPSTTL


TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



Plasmid Map:


ACCN: NM_001285506

ORF Size: 2157 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001285506.1](#), [NP_001272435.1](#)

RefSeq Size: 3002 bp

RefSeq ORF: 2160 bp

| | |
|----------------------|---|
| Locus ID: | 20111 |
| Cytogenetics: | 4 D2.3 |
| MW: | 81.8 kDa |
| Gene Summary: | <p>Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1) signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro-apoptotic function of BAD and DAPK1. In fibroblast, is required for EGF-stimulated phosphorylation of CREB1, which results in the subsequent transcriptional activation of several immediate-early genes. In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP. Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity. Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the pre-initiation complex. In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation. Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin-sensitive signaling independently of the PI3K/AKT pathway. Mediates cell survival by phosphorylating the pro-apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function. Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4). Mediates induction of hepatocyte proliferation by TGFA through phosphorylation of CEBPB (PubMed:10635333). Is involved in cell cycle regulation by phosphorylating the CDK inhibitor CDKN1B, which promotes CDKN1B association with 14-3-3 proteins and prevents its translocation to the nucleus and inhibition of G1 progression (By similarity). Phosphorylates EPHA2 at 'Ser-897', the RPS6KA-EPHA2 signaling pathway controls cell migration (By similarity).[UniProtKB/Swiss-Prot Function]</p> |