

## Product datasheet for MR231547

### Tek (NM\_001290551) Mouse Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Tek (NM\_001290551) Mouse Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** Tek  
**Synonyms:** AA517024; Cd202b; Hyk; STK1; Tie-2; Tie2  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**ORF Nucleotide Sequence:** >MR231547 representing NM\_001290551  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCCCGGATCGCC

ATGGACTCTTTAGCCGGCTTAGTTCTCTGTGGAGTCAGCTTGCTCCTTTATGGAGTAGTAGAAGGCGCCA  
 TGGACCTGATCTTGATCAATCCCTACCTCTTGCTGTGATGCCGAAACATCCCTCACCTGCATTGCCTC  
 TGGGTGGCACCCCATGAGCCCATCACCATAGGAAGGGACTTTGAAGCCTTAATGAACCAGCACCAAGAT  
 CCACTGGAGGTTACTCAAGATGTGACCAGAGAATGGGCGAAAAAGTTGTTTGAAGAGAGAAAAAGGCCA  
 GTAAGATTAATGGTGCTTATTTCTGTGAAGGTCGAGTTCCGAGGACAGGCTATAAGGATACGGACCATGAA  
 GATGCGTCAACAAGCATCCTTCCTACCTGCTACTTTAACTATGACCGTGGACAGGGGAGATAATGTGAAC  
 ATATCTTTCAAAAAGGTGTTAATTAAGAAGAAGATGCAGTGATTTACAAAAATGGATGTGAAGCTCAGA  
 AGTGGGGCCCGACTGTAGCCGTCCTTGACTACTTGCAAGAACAATGGAGTCTGCCATGAAGATACCGG  
 GGAATGCATTTGCCCTCCTGGGTTTATGGGGAGAACATGTGAGAAAGCTTGTGAGCCGCACACATTTGGC  
 AGGACCTGTAAGAAAAGGTGTAGTGGACCAGAAGGATGCAAGTCTTATGTGTTCTGTCTCCAGACCCTT  
 ACGGGTGTCTCTGTGCCACAGGCTGGAGGGGTTGCAGTGCAATGAAGCATGCCCATCTGGTTACTACGG  
 ACCAGACTGTAAGCTCAGGTGCCACTGTACCAATGAAGAGATATGTGATCGGTTCCAAGGATGCCTCTGC  
 TCTCAAGGATGGCAAGGGCTGCAGTGTGAGAAAAGGCAAGGCCAAGGATGACTCCACAGATAGAGGATT  
 TGCCAGATCACATTGAAGTAAACAGTGGAAAATTTAACCCTATCTGCAAAGCCTCTGGGTGGCCACTACC  
 TACTAGTGAAGAAATGACCCTAGTGAAGCCAGATGGGACAGTGTCCAACCAATGACTTCAACTATACA  
 GATCGTTTCTCAGTGGCCATATTCAGTGTCAACCGAGTCTTACCTCTGACTCAGGAGTCTGGGTCTGCA  
 GTGTGAACACAGTGGCTGGGATGGTGGAAAAGCCTTTCAACATTTCCGTCAAAGTTCTTCCAGAGCCCT  
 GCACGCCCAAAATGTGATTGACACTGGACATAACTTTGCTATCATCAATATCAGCTCTGAGCCTTACTTT  
 GGGGATGGACCCATCAAATCCAAGAAGCTTTTCTATAAACCTGTCAATCAGGCCTGGAAATACATTGAAG  
 TGACGAATGAGATTTTCACTCTCAACTACTTGGAGCCCGGACTGACTACGAGCTGTGTGTGCAGCTGGC  
 CCGTCTGGAGAGGGTGGAGAAGGGCATCTGGGCTGTGAGACGATTTACAACAGCGTCTATCGGACTC  
 CCTCTCCAAGAGGTCTCAGTCTCTGCCAAAAAGCCAGACAGCTCTAAATTTGACTTGGCAACCGATAT



[View online »](#)

TTACAAACTCAGAAGATGAATTTTATGTGGAAGTCGAGAGGCGATCCCTGCAAACAACAAGTGATCAGCA  
 GAACATCAAAGTGCCTGGGAACCTGACCTCGGTGCTACTGAGCAACTTAGTCCCCAGGGAGCAGTACACA  
 GTCCGAGCTAGAGTCAACACCAAGGCGCAGGGGAGTGGAGTGAAGAAGTCAAGGCTGGACCCTTAGTG  
 ACATTCTCCCTCAACAGAAAACATCAAGATCTCCAACATCACTGACTCCACAGCTATGGTTTCTTG  
 GACAATAGTGGATGGCTATTCGATTTCTCCATCATCATCCGGTATAAGGTTCAAGGCAAAAATGAAGAC  
 CAGCACATTGATGTGAAGATCAAGAATGCTACCGTTACTCAGTACCAGCTCAAGGCTCAGAGCCAGAGA  
 CTACATACCATGTGGATATTTTTGCTGAGAACAACATAGGATCAAGCAACCCAGCCTTTTCTCATGAAT  
 GAGGACGCTTCCACATTCAGCCTCTGCAGACCTCGGAGGGGAAAGATGCTACTCATAGCCATCCTT  
 GGGTCCGCTGGAATGACTTGCATCACCGTGTGTTGGCGTTTCTGATTATGTTGCAACTGAAGAGAGCAA  
 ATGTCCAAAGGAGAATGGCTCAGGCATTCCAGAACGTGAGAGAAGAACCAGCTGTGCAGTTAACTCAGG  
 AACTCTGGCCCTAACAGGAAGGCCAAAAACAATCCGGATCCCACAATTTATCCTGTGCTGACTGGAAT  
 GACATCAAGTTTCAAGACGTGATCGGAGAGGGCAACTTTGGCCAGGTTCTGAAGGCACGCATCAAGAAGG  
 ATGGGTTACGGATGGATGCCCCATCAAGAGGATGAAAGAGTATGCCTCAAAGATGATCACAGGGACTT  
 CGCAGGAGAACTGGAGGTTCTTTGTAACCTTGACACCATCCAACATCATTAACTCTTGGGAGCATGT  
 GAACACCGAGGCTATTTGTACCTAGCTATTGAGTATGCCCGCATGGAAACCTCTGGACTTCTCGCGTA  
 AGAGCAGAGTGTAGAGACAGACCTGCTTTTGCATCGCCAACAGTACAGCTTCCACACTGTCTCCCA  
 ACAGCTTCTTATTTGCTGCAGATGTGGCCCGGGGATGGACTACTTGAGCCAGAAACAGTTTATCCAC  
 AGGGACCTGGCTGCCAGAAAATTTTAGTTGGTAAAACCTACATAGCCAAAATAGCAGATTTTGGATTGT  
 CACGAGGTCAAGAAGTGTATGTAAAAAGACAATGGGAAGGCTCCCAGTGCCTGGATGGCAATCGAATC  
 ACTGAACATAGTGTCTATAACAACAACAGTGTGTCTGGTCTATGGTGTATTGCTCTGGGAGATTGTT  
 AGCTTAGGAGGCCACCCCTACTGCGGCATGACGTGCGCGGAGCTCTATGAGAAGTACCCAGGGCTACA  
 GGCTGGAGAAGCCCTGAACTGTGATGATGAGGTGTATGATCTAATGAGACAGTGTGGAGGGAGAAGCC  
 TTATGAGAGACCATCATTGCCCCAGATATTGGTGTCTTAAACAGGATGCTGGAAGAACGGAAGACATAC  
 GTGAACACCACACTGTATGAGAAGTTTACCTATGCAGGAATTGACTGCTCTGCGGAAGAAGCAGCC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**

>MR231547 representing NM\_001290551  
 Red=Cloning site Green=Tags(s)

MDSL AGLVLCGVSLLLYGVVEGAMDILINSLPLVSDAETSLTCIASGWHPHEPITIGRDFEALMNQHQD  
 PLEVTDVDTREWAKKVVWKREKASKINGAYFCEGRVVRGQAIRIRTMKMRQASFLPATLTMTVDRGDNVN  
 ISFKKVLIKEEDAVIYKNGCEAQKWGPDCSRPCTTCKNNGVCHEDTGEICPPGFMGRTCEKACEPHTFG  
 RTCKERCSGPEGCKSYVFLPDPYGCSCATGWRGLQCNEACPSGYYGPDCKLRCHCTNEEICDRFQGLC  
 SQGWQGLQCEKEGRPRMTPQIEDLPDHIEVNSGKFNPIKASGWPLPTSEEMTLVKPDGTVLQPNDFNYT  
 DRFSVAIFTVNRVLPDPSGVVWCSVNTVAGMVEKPFNISVKVLEPLHAPNVIDTGHNFAINISSEPYF  
 GDGPIKSKKLFYKPVNQAWKYIEVTNEIFTLNYLEPRTDYELCVQLARPGEGGEGHPGPVRRFTTASIGL  
 PPRGLSLLPKSQTALNLWQPIFTNSEDEFYVEVERSLQTTSDQQNIKVPGNLTSVLLSNLVPREQYT  
 VRRVNTKAQGEWSEELRAWLSDILPPQENIKISNITDSTAMVSWTIVDGYSISSIIIRYKVQGNED  
 QHIDVKIKNATVTQYQLKLEPETTYHVDIFAENNIGSSNPAFSHELRTLPHSPASADLGGGKMLLIAIL  
 GSAGMTCITVLLAFLIMLQLKRANVQRRMAQAFQNVREEPAVQFNSGTLALNRKAKNPDPTIYPVLDWN  
 DIKFQDVIGENFGQVLKARIKKDGLRMDAAIKRMKEYASKDDHRDFAGELEVLCCKLGHHPNIIINLLGAC  
 EHRGYLYLAIEYAPHGNLLDFLRKSRVLETDPAFAIANSTASTLSSQQLLHFAADVARGMDYLSQKQFIH  
 RDLAARNILVGENYIAKIADFGLSRGQEVYVVKTMGRLPVRWMAIESLNYSVYTTNSDVWSYGVLLWEIV  
 SLGGTPYCGMTCAELYEKLPQGYRLEKPLNCDDEVYDLMRQCWREKPYERPSFAQILVSLNRMLEERKTY  
 VNTTLYEKFTYAGIDCSAEAAA

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV


**Restriction Sites:**

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM\_001290551

ORF Size: 3216 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_001290551.1, NP_001277480.1</u>
<b>RefSeq Size:</b>	4549 bp
<b>RefSeq ORF:</b>	3219 bp
<b>Locus ID:</b>	21687
<b>Cytogenetics:</b>	4 43.34 cM
<b>MW:</b>	120.5 kDa
<b>Gene Summary:</b>	<p>Tyrosine-protein kinase that acts as cell-surface receptor for ANGPT1, ANGPT2 and ANGPT4 and regulates angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. Has anti-inflammatory effects by preventing the leakage of proinflammatory plasma proteins and leukocytes from blood vessels. Required for normal angiogenesis and heart development during embryogenesis. Required for post-natal hematopoiesis. After birth, activates or inhibits angiogenesis, depending on the context. Inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In quiescent vessels, ANGPT1 oligomers recruit TEK to cell-cell contacts, forming complexes with TEK molecules from adjoining cells, and this leads to preferential activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascades. In migrating endothelial cells that lack cell-cell adhesions, ANGPT1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1, and ultimately to the stimulation of sprouting angiogenesis. ANGPT1 signaling triggers receptor dimerization and autophosphorylation at specific tyrosine residues that then serve as binding sites for scaffold proteins and effectors. Signaling is modulated by ANGPT2 that has lower affinity for TEK, can promote TEK autophosphorylation in the absence of ANGPT1, but inhibits ANGPT1-mediated signaling by competing for the same binding site. Signaling is also modulated by formation of heterodimers with TIE1, and by proteolytic processing that gives rise to a soluble TEK extracellular domain. The soluble extracellular domain modulates signaling by functioning as decoy receptor for angiopoietins. TEK phosphorylates DOK2, GRB7, GRB14, PIK3R1, SHC1 and TIE1.[UniProtKB/Swiss-Prot Function]</p>