

Product datasheet for MR231611

Jak2 (NM_008413) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Jak2 (NM_008413) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Jak2
Synonyms:	Fd17
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR231611 representing NM_008413 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGAATGGCCTGCCTTACAATGACAGAAATGGAGGCAACCTCCACATCTCCTGTACATCAGAATGGTG
ATATTCCTGGAAGTGCTAATTCTGTGAAGCAGATAGAGCCAGTCCTTCAAGTGTATCTGTACCATTCTCT
TGGGCAAGCTGAAGGAGAGTATCTGAAGTTTCCAAGTGGAGAGTATGTTGCAGAAGAAATTTGTGTGGCT
GCTTCTAAAGCTTGTGGTATTACGCCTGTGTATCATAATATGTTTGCCTAATGAGTGAACCGAAAGGA
TCTGGTACCCACCAATCATGTCTTCCACATAGACGAGTCAACCAGGCATGACATACTCTACAGGATAAG
GTTCTACTTCCCTCATTGGTACTGTAGTGGCAGCAGCAGAACCTACAGATACGGAGTGTCCCGTGGGGCT
GAAGCTCCTCTGCTTGATGACTTTGTCTGCTTACCTTTTTGCTCAGTGGCGGCATGATTTTGTTCACG
GATGGATAAAAAGTACCTGTGACTCATGAAACTCAGGAAGAGTGTCTTGGGATGGCGGTGTAGACATGAT
GAGAATAGCTAAGGAGAAAGACCAGACTCCACTGGCTGTCTATAACTCTGTGAGCTACAAGACATTTCTTA
CCAAAGTGCCTTCGAGCGAAGATCCAAGACTATCACATTTAACC CGGAAGCGAATCAGGTACAGATTTCT
GCAGATTCATTGCAATTCAGTCAATGTAAGCCACTGCCAGGAACCTAAAACCTAAGTATCTTATAAA
CCTGGAACCCCTGCAGTCTGCCTTCTACACAGAACAGTTTGAAGTAAAAGAATCTGCAAGAGGTCCCTCA
GGTGAGGAGATTTTGAACCAATATAAATAACTGGAAACGGTGGAAATTCAGTGGTCAAGAGGGAACATA
AGGAAAGTGAGACTGACAGAACAGGACGTACAGTTATATTGTGATTTCCCTGATATTATTGATGTCAG
TATTAAGCAAGCAAACCAGGAATGCTCAAATGAAAGTAGAATTGTAAGTGTCCATAAACAAGATGGTAAA
GTTTTGGAGATAGAACTTAGCTCATTAAAAGAAGCCTTGTATTGTCGTCATTAATTGACGGGTATTACA
GACTAACTGCGGATGCGCACCAATTACCTCTGCAAAGAGGTGGCTCCCCAGCTGTGCTCGAGAACATACA
CAGCAACTGCCACGGCCCAATATCAATGGATTTTGCCATTAGCAAACCTAAAGAAGCGGGTAACCAGACT
GGACTATATGTGCTACGATGCAGCCCTAAGGACTTCAACAATACTTTCTGACCTTTGCTGTTGAGCGAG
AAAATGTCATTGAATATAAACTGTTTATTACGAAGAATGAGAATGGAGAATACAACCTCAGCGGGAC
TAAGAGGAACTCAGTAACCTTAAGGACCTTTGAATTGCTACCAGATGGAACTGTGCGCTCAGACAGT



[View online »](#)

ATCATCTTCCAGTTTACCAAATGCTGCCCCCAAAGCCAAAAGATAAATCAAACCTTCTCGTCTTCAGAA
CAAATGGTATTTCTGATGTTTCAGATCTCACCAACATTACAGAGGCATAATAATGTGAATCAAATGGTGT
TCACAAAATCAGGAATGAAGATTTAATATTTAATGAAAGTCTTGGCCAAGTACTTTTACAAAAATTTT
AAAGGTGAAGAAGAGAAGTTGGAGATTATGGTCAACTGCACAAAACGGAAGTCTTTTGAAGTCTTAG
ATAAAGCACATAGGAACATTTCAGAGTCTTTCTCGAAGCAGCAAGCATGATGAGTCAGCTTTCTCACA
GCATTTGGTTTTGAATTATGGTGTCTGTGTCTGTGGAGAGGAGAACATTCTGGTTCAAGAATTTGTA
TTTGGTCACTGGATACATACCTGAAGAAGAACAAAAATCCATAAATATATTATGGAACTTGGAGTGG
CTAAGCAGTTGGCATGGGCCATGCATTTTCTAGAAGAAAAATCCCTTATTCATGGGAATGTGTGTCTAA
AAATATCCTGCTTATCAGAGAAGAAAAACAGGAGAACGGGGAACCCACCTTTCATCAAACCTTAGTGATCCT
GGCATTAGCATTACAGTTCTACCGAAGGACATTCTCAGGAGAGAATACCATGGGTACCTCCTGAATGCA
TTGAGAATCCTAAAAATCTCAATCTGGCAACAGACAAGTGGAGCTTCGGGACCACTCTGTGGGAGATCTG
CAGTGGAGGAGATAAGCCCTGAGTGTCTGGATTCTCAAAGAAAGCTGCAGTTCATGAAGATAAGCAT
CAGCTTCTGCACCCAAGTGGACAGAGTTAGCAAACCTTATAAATAATTGCATGGACTATGAGCCAGATT
TCAGGCCCTGCTTTCAGAGCTGTCATCCGTGATCTTAACAGCCTGTTTACTCCAGATTATGAACTACTAAC
AGAAAATGACATGCTACCAAACATGAGAATAGGTGCCCTAGGGTTTTCTGGTGTCTTTGAAGACAGGGAC
CCTACACAGTTTTGAAGAGAGACACTTGAAGTTTCTACAGCAGCTTGGCAAAGGTAACCTCGGGAGTGTGG
AGATGTGCCGCTATGACCCGCTGCAGGACAACACTGGCGAGGTGGTTCGCTGTGAAGAAATCCAGCACAG
CACTGAAGAGCACCTCCGAGACTTTGAGAGGGAGATCGAGATCCTGAAATCCTTGCAGCATGACAACATC
GTCAAGTACAAGGGAGTGTGCTACAGTGCGGGTTCGGCGCAACCTAAGATTAATTATGGAATATTTACCAT
ATGGAAGTTTACGAGACTATCTCCAAAAACATAAAGAACGGATAGATCACAAAAAATCTTCAATACAC
AACATATTGGTGGAAAAATGAGAACAGGGTTAAAATAGGAGACTTCGGATTAACCAAAGTCTTGCCGAGG
ACAAAAGAATACTACAAAGTAAAGGAGCCAGGGGAAAGCCCATATTCTGGTACGCACCTGAATCCTTGAC
GGAGAGCAAGTTTTCTGTGGCCTCAGATGTGTGGAGCTTTGGAGTGGTTCTATACGAACTTTTACATAC
ATCGAGAAGAGTAAAAGTCCACCCGTGGAATTTATGCGAATGATTGGCAATGATAAAACAAGGGCAAATGA
TTGTGTTCCATTTGATAGAGCTACTGAAGAGCAACGGAAGATTGCCAAGGCCAGAAGGATGCCAGATGA
GATTTATGTGATCATGACAGAGTGTGGAACAACAATGTGAGCCAGCGTCCCTCCTTCAGGGACCTTTCC
CTTCGGGTGGATCAAATCCGGGACAGTATAGCTGCG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA


Protein Sequence: >MR231611 representing NM_008413
 Red=Cloning site Green=Tags(s)

MGMACLTMEATSTSPVHQNGDIPGSANSVKQIEPVLQVYLYHSLGQAEGEYLKFPSGEYVAEEICVA
 ASKACGITPVYHNMFMSETERIWYPPNHVFHIDESTRDILYRIRFYFPHWYCSGSSRTYRYGVSRGA
 EAPLLDDFVMSYLFQWRHDFVHGWIQVPVTHEEQEELGMAVLDMRIAKEKDQTPLAVYNSVSYKTFL
 PKCVRAIKIQDYHILTRKRIRYFRFRFIQQFSQCKATARNLKLKYLINLETLQSAFYTEQFEVKESARGPS
 GEEIFATIIITGNGGIQWSRGKHKESETLTEQDVQLYCDFPDIIDVSIKQANQECNESRIVTVHKQDGK
 VLEIELSSLKEALSFVSLIDGYRILTADAHYLCKEVAPPVLENIHNSNCHGPI SMDFAISKLKKAGNQT
 GLYVLRCPKDFNKYFLTF AVERENVIEYKHCLITKNENGEYNLSGTRKRFNLKDLLNCYQMETVRSDS
 IIFQFTKCCPPKPKDKSNLLVFRNGISDVQISPTLQRHNNVQMVFHKIRNEDLIFNESLGQGTFTKIF
 KGVRRVGDYQGLHKTEVLLKVLDAHRNYSSEFFEAASMSQLSHKHLVLNYGVCVCGEENILVQEFVK
 FGLSDTYLKKNKNSINILWKLGVAKQLAWAMHFLEEKSLIHGNVCAKNILLIREENRRTGNPPFIKLSDP
 GISITVLPKDILQERIPWVPECIENPKNLNLATDKWSFGTTLWEICSGGDKPLSALDSQRKLQFYEDKH
 QLPAPKWTELANLINNCOMDYEPDFRPAFRAVIRDLNSLFTPDYELLTENDMLPNMRIGALGFSGAFEDRD
 PTQFEERHLKFLQQLGKGNFGSVECRYDPLQDNTGEVAVKQLQHSTEEHLRDFEREIEILKSLQHDNI
 VKYKGVCSYAGRRNLRIMEYLPYGLRDYLVQKHKERIDHKLLQYTSQICKGMEYLGTKRYIHRDLATR
 NILVENENRVKIGDFGLTKVLPQDKEYYKVEKPESPIFWYAPESL TESKF SVASDVWSFGVVL YELFTY
 IEKSKSPVVEFMRMIGNDKQGMIVFHLIELLKSNGRLPRPEGCPDEIYVIMTECWNNNSQRPSFRDLS
 LRVDQIRDSIAA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI
Cloning Scheme:

Cloning sites used for ORF Shuttling:




* The last codon before the Stop codon of the ORF

ACCN: NM_008413
ORF Size: 3396 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_008413.2 , NM_008413.3 , NP_032439.2
RefSeq Size:	5055 bp
RefSeq ORF:	3399 bp
Locus ID:	16452
UniProt ID:	Q62120
Cytogenetics:	19 23.73 cM
MW:	130.6 kDa

Gene Summary:

Non-receptor tyrosine kinase involved in various processes such as cell growth, development, differentiation or histone modifications. Mediates essential signaling events in both innate and adaptive immunity. In the cytoplasm, plays a pivotal role in signal transduction via its association with type I receptors such as growth hormone (GHR), prolactin (PRLR), leptin (LEPR), erythropoietin (EPOR), thrombopoietin (THPO); or type II receptors including IFN-alpha, IFN-beta, IFN-gamma and multiple interleukins. Following ligand-binding to cell surface receptors, phosphorylates specific tyrosine residues on the cytoplasmic tails of the receptor, creating docking sites for STATs proteins. Subsequently, phosphorylates the STATs proteins once they are recruited to the receptor. Phosphorylated STATs then form homodimer or heterodimers and translocate to the nucleus to activate gene transcription. For example, cell stimulation with erythropoietin (EPO) during erythropoiesis leads to JAK2 autophosphorylation, activation, and its association with erythropoietin receptor (EPOR) that becomes phosphorylated in its cytoplasmic domain. Then, STAT5 (STAT5A or STAT5B) is recruited, phosphorylated and activated by JAK2. Once activated, dimerized STAT5 translocates into the nucleus and promotes the transcription of several essential genes involved in the modulation of erythropoiesis. Part of a signaling cascade that is activated by increased cellular retinol and that leads to the activation of STAT5 (STAT5A or STAT5B). In addition, JAK2 mediates angiotensin-2-induced ARHGEF1 phosphorylation. Plays a role in cell cycle by phosphorylating CDKN1B. Cooperates with TEC through reciprocal phosphorylation to mediate cytokine-driven activation of FOS transcription. In the nucleus, plays a key role in chromatin by specifically mediating phosphorylation of 'Tyr-41' of histone H3 (H3Y41ph), a specific tag that promotes exclusion of CBX5 (HP1 alpha) from chromatin.[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for MR231611