



ACTACCACCCAAAGATCAGGATCATCACTCAGATGCTGCAGTATCACAACAAGGCCCATCTGCTCAACAT  
 CCCCAGCTGGAAGTGGAAAGAGGGTGTGACGCAATATGCCTTGACAGCTCAAGTTGGGTTTCATAGCC  
 CAGAGCTGTCTGGCTCAAGGCCTCTCCACAATGCTTGCCAATCTCTTCTATGAGGTCATTCATAAAGA  
 TTGAGGAAGACACATGGCAGAAATACTACTTGGAAAGAGTCTCCAATGAAATGTACACAGAATATCTCTC  
 CAGTGCCTTCGTGGGTCTGTCTTCCCTACTGTTTGTGAGCTGTGTTTGTGAAGCTTAAGCTCCTGATG  
 ATAGCCATTGAGTACAAGTCTGCCAACAGAGAGAGCCGAAGCCGAAAGCGAATATTAATTAACCTGGGA  
 ACCACCTTAAGATCCAAGAAGTACTTTAGGATTTTTTCATCGCAAGTATGCCAAAGAAGTTAAAAGGGC  
 ATTTTTTACTGCAAGGCCTGTATGATGACGTACAGATCCCAAAGAATTAATAAATGTGGCTGCAGG  
 CGGCTCAAGGTTGAAGCTAGAGCCCGCTATCACAAAGACCCATTTATGCACAAGAATGCGACTCCCAATT  
 CTCCACACGTGCCAAGCCAGTTGAAGATGAGCAGCCGCAACCTGTACCAAAAAAAAAACAACGTAA  
 TGGGGGATGAGGAAGTCCGCAACACCTCCCGAAGCTGATGAGGCATGACCCCTTGTAAATTCCTGGC  
 AATGATCAGATTGACAACATGGACTCCAATGTAAAAAGTACGACTCCACTGGAATGTTTACTGGTGTG  
 CACCAAGGAGATTGAGAAAGTCATCTTACTCGAAGTGAAGTGCCTGACTGTCTGAGTGGCCATGT  
 CGTAGTCTGCATCTTGGGGATGTCAGCTCAGCCCTGATTGGCCTCCGGAACCTGGTATGCCACTTCGT  
 GCTAGCAACTTTCACTATCATGAGCTCAAACACATTGTGTTTGTGGGCTCCATTGAGTACCTCAAGAGGG  
 AGTGGGAAACTGCACAACCTCCCGAAGTGTCCATATTGCCTGGTACACCATTAAGTCGGGCTGATTT  
 AAGGGCTGTCAACATCAACCTCTGTGACATGTGCGTTATCCTGTGAGCCAATCAGAATAATATTGATGAT  
 ACTTCGCTTCCAGGACAAGGAATGCATCTTGGCGTCACTCAACATCAAATCTATGCAGTTTGTGACAGCA  
 TCGGGGCTTGCAGGCTAATCCCAAGGATTACACCTCCTGGAATGGACAGATCATACCCGACAACAG  
 CCCAGTGCACGGGATGTTACGCCAGCCGTCCATCACAACCTGGGGTCAACATTCATCATCAGGAACTC  
 GTGAATGATACCAATGTTTCAGTTTTTGGACCAAGACGATGACGATGACCTGACACAGAGCTGTACCTCA  
 CACAGCCCTTTGCTTGTGGGACAGCATTGCGGTGACGCTCCTGGACTCACTCATGAGCGCAGACTACTT  
 CAATGACAATATCCTCACCTAATACGGACCCTGGTGACAGGAGGAGCCACACCAGAGCTCGAGGCTCTA  
 ATAGCTGAGGAGAATGCACCTTCGAGGAGGCTACAGCACTCCGCAGACATTGGCCAACAGGACCGTTGCC  
 GAGTGGCCAGTTAGCCCTGTTAGATGGTCCCTTTCAGACTTAGGGGATGGTGGTTGTTATGGTATCT  
 GTTCTGCAAAGCTCTGAAAACATATAATATGCTTTGTTTTGGAATTTACCGGCTGAGAGATGCCACCTC  
 AGCACCCAGCCAGTGTACAAAAGGTACGTATCACCATCCTCCCTACGAGTTTGTGCTCGTACCAA  
 CAGACCTGATCTTCTGCCTGATGCAGTTTGACCACAACGCTGGCCAATCCCGGGCCAGTCTGTCTCATT  
 CTCCACTCCTCACAGTCGTCCAGTAAGAAGAGCTCCTCCGTCCACTCCATCCCGTCCACAGCAAATCGG  
 CCGAACCGGCCAAGTCCAGGGAGTCCCGCGACAAACAGAAGTACGTTTCAGGAAGAGCGGCTT

AGCGGACCGACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
 TGGATTACAAGGATGACGACGATAAGGTTTAA


**Protein Sequence:** >MR231689 representing NM\_001253368  
 Red=Cloning site Green=Tags(s)

MANGGGGGGGSSGGGGGGGGSGLRMSSNIHANHLSLDASSSSSSSSSSSSSSSSSSSSSVHEPKMDALI  
 IPVTMEVPCDSRQRMWAWFLASSMVTFFGGFLIILLWRTLKYLWTVCCCHCGGKTKEAQKINNGSSQADG  
 TLKPVDEKEEVAAEVGWMTSVKDWAGVMISAQTLTGRVLLVLFALSIGALVIYFIDSSNPIESCQNFY  
 KDFTLQIDMAFNVFLLYFGLRFIAANDKLWFLEVNVSVDFFTVPPVFVSVYLNRSWGLRFLRALRLI  
 QFSEILQFLNILKTSNSIKLVNLLSIFISTWLTAAAGFIHLVENSQDPWENFQNNQALTYWECVYLLMVTM  
 STVGYGDVYAKTTLGRLFMVFFILGGLAMFASYVPEIIEIIGNRKKYGGYSYSAVSGRKHIVVCGHITLES  
 VSNFLKDFLHKDRDDVNVEIVFLHNISPNELEALFKRHFTQVEFYQGSVLNPHDLARVKIESADACLIL  
 ANKYCADPDAEDASNIMRVISIKNYHPKIRIITQMLQYHNKAHLLNIPSWNKEGDDAICLAELKLGFA  
 QSCLAQGLSTMLANLFSMRSFIKIEEDTWQKYYLEGVSNEMYTEYLSSAFVGLSFPTVCELCFVKLKLMM  
 IAIEYKANSRERSRKRILINPGNHLKIQEGLGFFIASDAKEVKRAFFYCKACHDDVTPDKRIKCGCR  
 RLKVEARARYHKDPFMHKNATPNSPHVPKPVEDEQPPTLSPKKKQRNGGMRNSPNTSPKLMRHDPLLI  
 NDQIDNMDSNVKKYDSTGMFHCAPKEIEKVILTRSEAAMTVLSGHVVVCFGDVSSALIGLRNLVMP  
 LRSNFHYHELKHIIVFVGSIEYLKREWETLHNFPKVSILPGTPLSRADLRAVINLDCMVCVILSANQNNID  
 TSLQDKECILASLNIKSMQFDDSIGVLQANSQGFPPGMDRSPDNPVHGMLRQPSITTVGNIPITEL  
 VNDTNVQFLDQDDDDPDTELYLTQPFACGTAFVSVLDSLMSATYFNDNILLIRTLVTGGATPELEAL  
 IAEENALRGGYSTPQTLANRDRCRVAQLALLDGPFDLGDGGCYGDLFCALKTYNMLCFGIYRLRDAHL  
 STPSQCTKRYVITNPPYEFELVPTDLIFCLMQFDHNAQSRASLSHSSHSQSSSKSSSVHSIPSTANR  
 PNRPKSRESRDKQKYVQEERL


SGP TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

**Restriction Sites:** SgfI-RsrII

**Cloning Scheme:**



\* The last codon before the Stop codon of the ORF

**Plasmid Map:**


**ACCN:** NM\_001253368

**ORF Size:** 3633 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001253368.1](#), [NP\\_001240297.1](#)

**RefSeq Size:** 6119 bp

**RefSeq ORF:** 3636 bp

**Locus ID:** 16531

**Cytogenetics:** 14 A3

**MW:** 135.3 kDa

**Gene Summary:**

Potassium channel activated by both membrane depolarization or increase in cytosolic Ca(2+) that mediates export of K(+). It is also activated by the concentration of cytosolic Mg(2+). Its activation dampens the excitatory events that elevate the cytosolic Ca(2+) concentration and/or depolarize the cell membrane. It therefore contributes to repolarization of the membrane potential. Plays a key role in controlling excitability in a number of systems, such as regulation of the contraction of smooth muscle, the tuning of hair cells in the cochlea, regulation of transmitter release, and innate immunity. In smooth muscles, its activation by high level of Ca(2+), caused by ryanodine receptors in the sarcoplasmic reticulum, regulates the membrane potential. In cochlea cells, its number and kinetic properties partly determine the characteristic frequency of each hair cell and thereby helps to establish a tonotopic map. Kinetics of KCNMA1 channels are determined by alternative splicing, phosphorylation status and its combination with modulating beta subunits. Highly sensitive to both iberiotoxin (IbTx) and charybdotoxin (CTX).[UniProtKB/Swiss-Prot Function]