

ACTACCACCCAAAGATCAGGATCATCACTCAGATGCTGCAGTATCACAACAAGGCCCATCTGCTCAACAT
CCCCAGCTGGAAGTGGAAAGAGGGTGTGACGCAATATGCCTTGACAGCTCAAGTTGGGTTTCATAGCC
CAGAGCTGTCTGGCTCAAGGCCTCTCCACAATGCTTGCCAATCTCTTCTATGAGGTCATTCAAAGA
TTGAGGAAGACACATGGCAGAAATACTACTTGGAAAGAGTCTCCAATGAAATGTACACAGAATATCTCTC
CAGTGCCTTCGTGGGTCTGTCTCCCTACTGTTTGTGAGCTGTGTTTGTGAAGCTTAAGCTCCTGATG
ATAGCCATTGAGTACAAGTCTGCCAACAGAGAGAGCCGAATATTAATTAACCCGGGAACCACCTTAAGA
TCCAAGAAGGTACTTTAGGATTTTTATCGCAAGTGTGCCAAAGAAGTTAAAAGGGCATTTTTTTACTG
CAAGGCCTGCATGATGACGTACAGATCCCAAAAGAATTAAAAAATGTGGCTGCAGGCGCCCAAGATG
TCCATCTACAAGAGAATGAGACGAGCATGTTGTTTTGATTGCGGACGTTCTGAGCGTGACTGCTCGTGCA
TGTCAGGCCGTGTGCGTGGTAACGTGGACACCCTTGAGAGAACCCTTCCCGCTTTCTTCTGTCTGTAA
TGATTGCTCCACCAGTTTCCGTGCCTTTGAAGATGAGCAGCCGCAACCCTGTCACCAAAAAAAAAACAA
CGTAATGGGGCATGAGGAACCTGCCAACACCTCCCCGAAGCTGATGAGGCATGACCCCTTGTAAATTC
CTGGCAATGATCAGATTGACAACATGGACTCCAATGTGAAAAAGTACGACTCCACTGGAATGTTTACTG
GTGTGCACCCAAGGAGATTGAGAAAGTCATCTTGACTCGAAGTGAAGCTGCCATGACTGTCTGAGTGGC
CATGTCTGATGCTGATCTTTGGGGATGTCAGCTCAGCCCTGATTGGCCTCCGGAACCTGGTGATGCCAC
TTGCTGTAGCAACTTTCACTATCATGAGCTCAAACACATTGTGTTTGTGGGCTCCATTGAGTACCTCAA
GAGGGAGTGGGAAACTGCACAACCTCCCGAAAGTGTCCATATTGCCTGGTACACCATTAAGTCGGGCT
GATTTAAGGGCTGTCAACATCAACCTCTGTGACATGTGCGTTATCCTGTGACCCAATCAGAATAATTTG
ATGATACTTCGCTTCAGGACAAGGAATGCATCTTGGCGTCACTCAACATCAAAATCTATGCAAGTTGATGA
CAGCATCGGGTCTTGCAGGCTAATCCCAAGGATTACACCTCCTGGAATGGACAGATCATCACCCGAC
AACAGCCAGTGCACGGGATGTTACGCCAGCCGTCCATCACAACCTGGGGTCAACATTCATCATACGG
AACTCGTGAATGATACCAATGTTCAAGTTTTGGACCAAGACGATGACGATGACCCTGACACAGAGCTGTA
CCTCACACAGCCCTTTGCTTGTGGGACAGCATTGCGCTCAGCGTCTGGACTCACTCATGAGCGCGACA
TACTTCAATGACAATATCCTCACCTAATACGGACCCTGGTGACAGGAGGAGCCACACCAGAGCTCGAGG
CTCTAATAGCTGAGGAGAATGCACCTCGAGGAGCTACAGCACTCCGCAGACATTGGCCAACAGGACCG
TTGCCGAGTGGCCAGTTAGCCCTGTTAGATGGTCCCTTTCAGACTTAGGGGATGGTGGTTGTTATGGT
GATCTGTTCTGCAAAGCTCTGAAAACATATAATATGCTTTGTTTTGGAATTTACCGGCTGAGAGATGCC
ACCTCAGCACCCAGCCAGTGTACAAAAGGTACGTATCACCATCCTCCCTACGAGTTTGTGCTCGT
ACCAACAGACCTGATCTTCTGCCTGATGCAGTTTGACCACAACGCTGGCCAATCCCGGGCCAGTCTGTCT
CATTCTCCCACTCCTCACAGTGTCCAGTAAGAAGAGCTCCTCCGTCCACTCCATCCCGTCCACAGCAA
ATCGGCCGAACCGGCCAAGTCCAGGGAGTCCCGGACAAACAGAAGTACGTTTACGGAAGAGCGGCTT

AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
TGGATTACAAGGATGACGACGATAAGGTTAA


Protein Sequence: >MR231714 representing NM_001253364
 Red=Cloning site Green=Tags(s)

MANGGGGGGGSSGGGGGGGGSGLRMSSNIHANHLSLDASSSSSSSSSSSSSSSSSSSSSVHEPKMDALI
 IPVTMEVPCDSRGQRMWVAFLASSMVTFFGGFLIILLWRTLKYLWTVCCCHCGGKTKEAQKINNGSSQADG
 TLKPVDEKEEVAAEVGWMTSVKDWAGVMISAQTLTGRVLLVLFALSIGALVIYFIDSSNPIESCQNFY
 KDFTLQIDMAFNVFLLYFGLRFIAANDKLWFLEVNVSVDFFTVPPVFVSVYLNRSWGLRFLRALRLI
 QFSEILQFLNILKTSNSIKLVNLLSIFISTWLTAAAGFIHLVENSQDPWENFQNNQALTYWECVYLLMVTM
 STVGYGDVYAKTTLGRLFMVFFILGGLAMFASYVPEIIEIIGNRKKYGGYSYSAVSGRKHIVVCGHITLES
 VSNFLKDFLHKDRDDVNVEIVFLHNISPNELEALFKRHFTQVEFYQGSVLNPHDLARVKIESADAACLIL
 ANKYCADPDAEDASNIMRVISIKNYHPKIRIITQMLQYHNKAHLLNIPSWNKEGDDAICLAELKLGFAIA
 QSCLAQGLSTMLANLFSMRSFIKIEEDTWQKYYLEGVSNEMYTEYLSAFVGLSFPTVCELFCVKLKLML
 IAIEYKSNANRESRILINPGNHLKIQEGTLGFFIASDAKEVKRAFFYCKACHDDVTDPKRIKCKGCRPPKM
 SIYKRMRRACCFDCGRSERDCSCMSGRVVRGNVDLERTFPLSSVSVNDCSTSFRAFEDEQPPTLSPKKKQ
 RNGGMRNSPNTSPKLMRHDPLLIPGNDQIDNMDSNVKKYDSTGMFHWCAPKEIEKVILTRSEAMTVLSG
 HVVVCIFGDVSSALIGLRNLVMPLRASNFHYHELKHIYVFGSIEYLKREWETLHNFPKVSIILPGTPLSRA
 DLRAVINLDCMCVILSANQNNIDDTSLQDKECILASLNIKSMQFDDSIGVLQANSQGFPPGMDRSSPD
 NSPVHGMRLRQPSITTVNIPITTELVDNTNVQFLDQDDDDDPTELYLTQPFACGTAFVSVLDSLMSAT
 YFNDNILTLIRTLVTGGATPELEALIAEENALRGGYSTPQTLANRDRCRVAQLALLDGPFDLGDGGCYG
 DLFCALKTYNMLCFGIYRLDAHLSTPSQCTKRYVITNPPYEFELVPTDLIFCLMQFDHNAGQSRASLS
 HSSHSSQSSSKSSSVHSIPSTANRPNRPKSRESRDKQKYVQEERL

SGP TRTRPLEQKLI SEEDLAANDILDYKDDDDKV


Restriction Sites: SgfI-RsrII

Cloning Scheme:



* The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001253364

ORF Size: 3708 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001253364.1](#), [NP_001240293.1](#)

RefSeq Size: 6194 bp

RefSeq ORF: 3711 bp

Locus ID: 16531

Cytogenetics: 14 A3

MW: 138 kDa

Gene Summary:

Potassium channel activated by both membrane depolarization or increase in cytosolic Ca(2+) that mediates export of K(+). It is also activated by the concentration of cytosolic Mg(2+). Its activation dampens the excitatory events that elevate the cytosolic Ca(2+) concentration and/or depolarize the cell membrane. It therefore contributes to repolarization of the membrane potential. Plays a key role in controlling excitability in a number of systems, such as regulation of the contraction of smooth muscle, the tuning of hair cells in the cochlea, regulation of transmitter release, and innate immunity. In smooth muscles, its activation by high level of Ca(2+), caused by ryanodine receptors in the sarcoplasmic reticulum, regulates the membrane potential. In cochlea cells, its number and kinetic properties partly determine the characteristic frequency of each hair cell and thereby helps to establish a tonotopic map. Kinetics of KCNMA1 channels are determined by alternative splicing, phosphorylation status and its combination with modulating beta subunits. Highly sensitive to both iberiotoxin (IbTx) and charybdotoxin (CTX).[UniProtKB/Swiss-Prot Function]