

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for PH300436

#### PSMD8 (NM\_002812) Human Mass Spec Standard

### Product data:


Product Type:	Mass Spec Standards
Description:	PSMD8 MS Standard C13 and N15-labeled recombinant protein (NP_002803)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC200436
Predicted MW:	30 kDa
Protein Sequence:	<pre>&gt;RC200436 protein sequence Red=Cloning site Green=Tags(s)</pre>
	MYEQLKGEWNRKSPNLSKCGEELGRLKLVLLELNFLPTTGTKLTKQQLILARDILEIGAQWSILRKDIPS FERYMAQLKCYYFDYKEQLPESAYMHQLLGLNLLFLLSQNRVAEFHTELERLPAKDIQTNVYIKHPVSLE QYLMEGSYNKVFLAKGNIPAESYTFFIDILLDTIRDEIAGCIEKAYEKILFTEATRILFFNTPKKMTDYA KKRGWVLGPNNYYSFASQQQKPEDTTIPSTELAKQVIEYARQLEMIV
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 $\mu$ g/ $\mu$ L as determined by microplate BCA method
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	<u>NP 002803</u>
RefSeq Size:	1556
RefSeq ORF:	771
Synonyms:	HEL-S-91n; HIP6; HYPF; Nin1p; p31; Rpn12; S14
Locus ID:	5714
UniProt ID:	<u>P48556, V9HW09</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PSMD8 (NM_002812) Human Mass Spec Standard – PH300436
Cytogenetics:	19q13.2
Summary:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 1. [provided by RefSeq, Jul 2008]
Protein Pathway	s: Proteasome

## **Product images:**



Coomassie blue staining of purified PSMD8 protein (Cat# [TP300436]). The protein was produced from HEK293T cells transfected with PSMD8 cDNA clone (Cat# [RC200436]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US