

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for PH301701

p38 (CRK) (NM_016823) Human Mass Spec Standard

Product data:

Product Type:	Mass Spec Standards	
Description:	CRK MS Standard C13 and N15-labeled recombinant protein (NP_058431)	
Species:	Human	
Expression Host:	HEK293	
Expression cDNA Clone or AA Sequence:	RC201701	
Predicted MW:	33.8 kDa	
Protein Sequence:	<pre>>RC201701 protein sequence Red=Cloning site Green=Tags(s)</pre>	
	MAGNFDSEERSSWYWGRLSRQEAVALLQGQRHGVFLVRDSSTSPGDYVLSVSENSRVSHYIINSSGPRPP VPPSPAQPPPGVSPSRLRIGDQEFDSLPALLEFYKIHYLDTTTLIEPVSRSRQGSGVILRQEEAEYVRAL FDFNGNDEEDLPFKKGDILRIRDKPEEQWWNAEDSEGKRGMIPVPYVEKYRPASASVSALIGGNQEGSHP QPLGGPEPGPYAQPSVNTPLPNLQNGPIYARVIQKRVPNAYDKTALALEVGELVKVTKINVSGQWEGECN GKRGHFPFTHVRLLDQQNPDEDFS	
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV	
Tag:	C-Myc/DDK	
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining	
Concentration:	>0.05 µg/µL as determined by microplate BCA method	
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine	
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3	
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.	
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.	
RefSeq:	<u>NP 058431</u>	
RefSeq Size:	3225	
RefSeq ORF:	912	
Synonyms:	CRKII; p38	
Locus ID:	1398	



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE p38 (CRK) (NM_016823) Human Mass Spec Standard – PH301701		
UniProt ID:	<u>P46108, A0A0S2Z3Q4, L7RT18</u>	
Cytogenetics:	17p13.3	
Summary:	This gene encodes a member of an adapter protein family that binds to several tyrosine- phosphorylated proteins. The product of this gene has several SH2 and SH3 domains (src- homology domains) and is involved in several signaling pathways, recruiting cytoplasmic proteins in the vicinity of tyrosine kinase through SH2-phosphotyrosine interaction. The N- terminal SH2 domain of this protein functions as a positive regulator of transformation whereas the C-terminal SH3 domain functions as a negative regulator of transformation. Two alternative transcripts encoding different isoforms with distinct biological activity have been described. [provided by RefSeq, Jul 2008]	
Protein Families:	Druggable Genome, Transcription Factors	
Protein Pathways:	Chemokine signaling pathway, Chronic myeloid leukemia, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways in cancer, Regulation of actin cytoskeleton, Renal cell carcinoma	

Product images:

116 —	-
66 —	-
45 —	-
35 —	-
25 —	
18 14	:

Coomassie blue staining of purified CRK protein (Cat# [TP301701]). The protein was produced from HEK293T cells transfected with CRK cDNA clone (Cat# [RC201701]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US