

## **Product datasheet for PH302459**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Chemokine Receptor D6 (ACKR2) (NM 001296) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** CCBP2 MS Standard C13 and N15-labeled recombinant protein (NP\_001287)

Species: Human **HEK293 Expression Host: Expression cDNA Clone** 

or AA Sequence:

RC202459

Predicted MW: 43.4 kDa

>RC202459 protein sequence **Protein Sequence:** 

Red=Cloning site Green=Tags(s)

MAATASPQPLATEDADSENSSFYYYDYLDEVAFMLCRKDAVVSFGKVFLPVFYSLIFVLGLSGNLLLLMV LLRYVPRRRMVEIYLLNLAISNLLFLVTLPFWGISVAWHWVFGSFLCKMVSTLYTINFYSGIFFISCMSL DKYLEIVHAQPYHRLRTRAKSLLLATIVWAVSLAVSIPDMVFVQTHENPKGVWNCHADFGGHGTIWKLFL RFQQNLLGFLLPLLAMIFFYSRIGCVLVRLRPAGQGRALKIAAALVVAFFVLWFPYNLTLFLHTLLDLQV FGNCEVSQHLDYALQVTESIAFLHCCFSPILYAFSSHRFRQYLKAFLAAVLGWHLAPGTAQASLSSCSES

SILTAQEEMTGMNDLGERQSENYPNKEDVGNKSA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

>0.05 µg/µL as determined by microplate BCA method **Concentration:** 

**Labeling Method:** Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 001287

RefSeg Size: 2978 RefSeq ORF: 1152

Synonyms: CCBP2; CCR9; CCR10; CMKBR9; D6; hD6

Locus ID: 1238





UniProt ID: <u>000590</u>, <u>A1LP82</u>

Cytogenetics: 3p22.1


**Summary:** This gene encodes a beta chemokine receptor, which is predicted to be a seven

transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptor-mediated signal transduction are critical for the recruitment of effector immune cells to the inflammation site. This gene is expressed in a range of tissues and hemopoietic cells. The expression of this receptor in lymphatic endothelial cells and overexpression in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and possible chemokine effects on the development and growth of vascular tumors. This receptor appears to bind the majority of beta-chemokine family members; however, its specific function remains unknown. This gene is mapped to chromosome 3p21.3, a region that

includes a cluster of chemokine receptor genes. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, GPCR, Transmembrane

## **Product images:**



Coomassie blue staining of purified ACKR2 protein (Cat# [TP302459]). The protein was produced from HEK293T cells transfected with ACKR2 cDNA clone (Cat# [RC202459]) using MegaTran 2.0 (Cat# [TT210002]).