

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for PH304709

Hemoglobin subunit gamma 2 (HBG2) (NM_000184) Human Mass Spec Standard

Product data:


Product Type:	Mass Spec Standards
Description:	HBG2 MS Standard C13 and N15-labeled recombinant protein (NP_000175)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC204709
Predicted MW:	16.1 kDa
Protein Sequence:	>RC204709 protein sequence Red=Cloning site Green=Tags(s)
	MGHFTEEDKATITSLWGKVNVEDAGGETLGRLLVVYPWTQRFFDSFGNLSSASAIMGNPKVKAHGKKVLT SLGDAIKHLDDLKGTFAQLSELHCDKLHVDPENFKLLGNVLVTVLAIHFGKEFTPEVQASWQKMVTAVAS ALSSRYH
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	<u>NP 000175</u>
RefSeq Size:	583
RefSeq ORF:	441
Synonyms:	HBG-T1; TNCY
Locus ID:	3048
UniProt ID:	<u>P69892, D9YZU9</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Hemoglobin subunit gamma 2 (HBG2) (NM_000184) Human Mass Spec Standard – PH304709
Cytogenetics:	11p15.4
Summary:	The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma product (HBG2) and alanine is found in the A-gamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'- epsilon gamma-G gamma-A delta beta3'. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified HBG2 protein (Cat# [TP304709]). The protein was produced from HEK293T cells transfected with HBG2 cDNA clone (Cat# [RC204709]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US