

Product datasheet for PH305752

UGT1A6 (NM_205862) Human Mass Spec Standard

Product data:

Product Type:	Mass Spec Standards
Description:	UGT1A6 MS Standard C13 and N15-labeled recombinant protein (NP_995584)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC205752
Predicted MW:	60.7 kDa
Protein Sequence:	>RC205752 protein sequence Red=Cloning site Green=Tags(s)

MACLLRSFQRISAGVFFLALWGMVVGDKLLVVPQDGSOWL SMKDI VEVLSDRGHEIVVVVPEVNLLKES
KYYTRKIYPVPYDQEELKNRYQSFSGNNHFAERSFLTAPQTEYRNNMIVIGLYFINCQSLQDRDTLNFFK
ESKFDALFTDPALPCGVILAEYGLPSVYLFRGFPCSLEHTFSRSPDPVSYIPRCYTKFSDHMTFSQRVA
NFLVNLLEPYLFYCLFSKYEELASAVLKRVDIITLYQKVSVWLLRYDFVLEYPRPVMPNMVFIGGINCK
KRKDLSEFEAYINASGEHGIVVFSLGSMVSEIPEKKAMATADALGKIPQTVLWRYTGTRPSNLANTIL
VKWLPQNDLLGHPMTRAFITHAGSHGVYESICNGVPMVMPLFGDQMDNAKRMETKGAGVTLNVLEMTSE
DLENALKAVINDKSYKENIMRLSSLHKDRPVEPLDLAVFWVEFVMRHKGAPHLRPAHADL TWYQYHSLDV
IGFLLAVVLTVAFITFKCCAYGRKCLGKKGRVKKAHKSKTH

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	<u>NP_995584</u>
RefSeq Size:	1691
RefSeq ORF:	1599



[View online »](#)

Synonyms:	GNT1; HLUGP; HLUGP1; hUG-BR1; UDPGT; UDPGT 1-6; UGT-1A; UGT-1C; UGT-1E; UGT-1F; UGT1; UGT1-01; UGT1-03; UGT1-05; UGT1-06; UGT1.1; UGT1.3; UGT1.5; UGT1.6; UGT1A; UGT1A1; UGT1A3; UGT1A5; UGT1A6S; UGT1C; UGT1E; UGT1F
Locus ID:	54578
UniProt ID:	Q5DT01
Cytogenetics:	2q37.1
Summary:	<p>This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The enzyme encoded by this gene is active on phenolic and planar compounds. Alternative splicing in the unique 5' end of this gene results in two transcript variants. [provided by RefSeq, Jul 2008]</p>
Protein Families:	Transmembrane
Protein Pathways:	Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

Product images:



Coomassie blue staining of purified UGT1A6 protein (Cat# [TP305752]). The protein was produced from HEK293T cells transfected with UGT1A6 cDNA clone (Cat# [RC205752]) using MegaTran 2.0 (Cat# [TT210002]).