

Product datasheet for PH310623

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Protein Kinase A regulatory subunit I alpha (PRKAR1A) (NM_212471) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: PRKAR1A MS Standard C13 and N15-labeled recombinant protein (NP_997636)

Species: Human

Expression Host: HEK293

Expression cDNA Clone RC210623

or AA Sequence:

Predicted MW: 43 kDa

Protein Sequence: >RC210623 protein sequence

Red=Cloning site Green=Tags(s)

MESGSTAASEEARSLRECELYVQKHNIQALLKDSIVQLCTARPERPMAFLREYFERLEKEEAKQIQNLQK AGTRTDSREDEISPPPPNPVVKGRRRRGAISAEVYTEEDAASYVRKVIPKDYKTMAALAKAIEKNVLFSH LDDNERSDIFDAMFSVSFIAGETVIQQGDEGDNFYVIDQGETDVYVNNEWATSVGEGGSFGELALIYGTP RAATVKAKTNVKLWGIDRDSYRRILMGSTLRKRKMYEEFLSKVSILESLDKWERLTVADALEPVQFEDGQ KIVVQGEPGDEFFIILEGSAAVLQRRSENEEFVEVGRLGPSDYFGEIALLMNRPRAATVVARGPLKCVKL

DRPRFERVLGPCSDILKRNIQQYNSFVSLSV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 997636

RefSeq Size: 4518 RefSeq ORF: 1143

Synonyms: ACRDYS1; ADOHR; CAR; CNC; CNC1; PKR1; PPNAD1; PRKAR1; TSE1





Protein Kinase A regulatory subunit I alpha (PRKAR1A) (NM_212471) Human Mass Spec Standard - PH310623

Locus ID: 5573

UniProt ID: <u>P10644</u>, <u>B2R5T5</u>

Cytogenetics: 17q24.2

Summary: cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its


effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. This gene encodes one of the regulatory subunits. This protein was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids. Mutations in this gene cause Carney complex (CNC). This gene can fuse to the RET protooncogene by gene rearrangement and form the thyroid tumor-specific chimeric oncogene known as PTC2. A nonconventional nuclear localization sequence (NLS) has been found for this protein which suggests a role in DNA replication via the protein serving as a nuclear transport protein for the second subunit of the Replication Factor C (RFC40). Several alternatively spliced transcript variants encoding

two different isoforms have been observed. [provided by RefSeq, Jan 2013]

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Apoptosis, Insulin signaling pathway

Product images:



Coomassie blue staining of purified PRKAR1A protein (Cat# [TP310623]). The protein was produced from HEK293T cells transfected with PRKAR1A cDNA clone (Cat# [RC210623]) using MegaTran 2.0 (Cat# [TT210002]).