

Product datasheet for PH310728

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP6V1F (NM 004231) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: ATP6V1F MS Standard C13 and N15-labeled recombinant protein (NP_004222)

Species:HumanExpression Host:HEK293

Expression cDNA Clone

RC210728

or AA Sequence:

Predicted MW: 13.4 kDa

Protein Sequence: >RC210728 protein sequence

Red=Cloning site Green=Tags(s)

MAGRGKLIAVIGDEDTVTGFLLGGIGELNKNRHPNFLVVEKDTTINEIEDTFRQFLNRDDIGIILINQYI

AEMVRHALDAHQQSIPAVLEIPSKEHPYDAAKDSILRRARGMFTAEDLR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 004222

RefSeq Size: 748 RefSeq ORF: 357

Synonyms: ATP6S14; VATF; Vma7

Locus ID: 9296

UniProt ID: Q16864, A4D1K0

Cytogenetics: 7q32.1






Summary:

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c", and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is the V1 domain F subunit protein. [provided by RefSeq, Jul 2008]

Protein Pathways:

Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

Product images:



Coomassie blue staining of purified ATP6V1F protein (Cat# [TP310728]). The protein was produced from HEK293T cells transfected with ATP6V1F cDNA clone (Cat# [RC210728]) using MegaTran 2.0 (Cat# [TT210002]).