

## **Product datasheet for PH314533**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CRYGB (NM 005210) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

Description: CRYGB MS Standard C13 and N15-labeled recombinant protein (NP\_005201)

Species: Human **HEK293 Expression Host:** 

**Expression cDNA Clone** 

RC214533

or AA Sequence: Predicted MW:

20.9 kDa

>RC214533 protein sequence **Protein Sequence:** 

Red=Cloning site Green=Tags(s)

MGKITFYEDRAFQGRSYECTTDCPNLQPYFSRCNSIRVESGCWMIYERPNYQGHQYFLRRGEYPDYQQWM GLSDSIRSCCLIPPHSGAYRMKIYDRDELRGQMSELTDDCLSVQDRFHLTEIHSLNVLEGSWILYEMPNY

RGRQYLLRPGEYRRFLDWGAPNAKVGSLRRVMDLY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Labeling Method:** Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

25 mM Tris-HCl, 100 mM glycine, pH 7.3 **Buffer:** 

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 005201

RefSeg Size: 643 RefSeq ORF: 525

Synonyms: CRYG2; CTRCT39

Locus ID: 1419 UniProt ID: P07316





Cytogenetics:

2q33.3


Summary:

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. Four gamma-crystallin genes (gamma-A through gamma-D) and three pseudogenes (gamma-E, gamma-F, gamma-G) are tandemly organized in a genomic segment as a gene cluster. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation. [provided by RefSeq, Jul 2008]

**Protein Families:** 

Druggable Genome

## **Product images:**



Coomassie blue staining of purified CRYGB protein (Cat# [TP314533]). The protein was produced from HEK293T cells transfected with CRYGB cDNA clone (Cat# [RC214533]) using MegaTran 2.0 (Cat# [TT210002]).