

## Product datasheet for PH315410

### RGS10 (NM\_002925) Human Mass Spec Standard

#### Product data:


Product Type:	Mass Spec Standards
Description:	RGS10 MS Standard C13 and N15-labeled recombinant protein (NP_002916)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC215410
Predicted MW:	19.4 kDa
Protein Sequence:	>RC215410 representing NM_002925 Red=Cloning site Green=Tags(s)  MEHIHSDGSSSSSHQSLKSTAKWAASLENLLEDPEGVKRFREFLKKEFSEENVLFWLACEDFKKMQDKT QMQEKAKEIYMTFLSSKASSQVNVGQSRLNEKILEEPHPLMFQKLQDQIFNLMKYDSYSRFLKSDLFLK HKRTEEEEEEDLPDAQTAAKRASRIYNT  TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- <sup>13</sup> C <sub>6</sub> , <sup>15</sup> N <sub>4</sub> ]-L-Arginine and [U- <sup>13</sup> C <sub>6</sub> , <sup>15</sup> N <sub>2</sub> ]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	<a href="#">NP_002916</a>
RefSeq Size:	859
RefSeq ORF:	501
Locus ID:	6001
UniProt ID:	<a href="#">O43665</a>
Cytogenetics:	10q26.11



[View online »](#)

**Summary:**

Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 10 belongs to this family. All RGS proteins share a conserved 120-amino acid sequence termed the RGS domain. This protein associates specifically with the activated forms of the two related G-protein subunits, G-alpha<sub>i3</sub> and G-alpha<sub>z</sub> but fails to interact with the structurally and functionally distinct G-alpha subunits. Regulator of G protein signaling 10 protein is localized in the nucleus. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Product images:**

Coomassie blue staining of purified RGS10 protein (Cat# [TP315410]). The protein was produced from HEK293T cells transfected with RGS10 cDNA clone (Cat# [RC215410]) using MegaTran 2.0 (Cat# [TT210002]).