

Product datasheet for PH315482

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PSMF1 (NM 178578) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: PSMF1 MS Standard C13 and N15-labeled recombinant protein (NP 848693)

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

RC215482

or AA Sequence: Predicted MW:

29.8 kDa

>RC215482 protein sequence **Protein Sequence:**

Red=Cloning site Green=Tags(s)

MAGLEVLFASAAPAITCRODALVCFLHWEVVTHGYCGLGVGDOPGPNDKKSELLPAGWNNNKDLYVLRYE YKDGSRKLLVKAITVESSMILNVLEYGSQQVADLTLNLDDYIDAEHLGDFHRTYKNSEELRSRIVSGIIT PIHEQWEKANVSSPHREFPPATAREVDPLRIPPHHPHTSRQPPWCDPLGPFVVGGEDLDPFGPRRGGMIV

DPLRSGFPRALIDPSSGLPNRLPPGAVPPGARFDPFGPIGTSPPGPNPDHLPPPGYDDMYL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

Labeling Method: Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 848693

RefSeq Size: 3686 RefSeq ORF: 813 Synonyms: PI31 9491 Locus ID:

UniProt ID: Q92530, A0A140VJT2





Cytogenetics:

20p13


Summary:

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a protein that inhibits the activation of the proteasome by the 11S and 19S regulators. Alternative transcript variants have been identified for this gene. [provided by RefSeq, Jul 2008]

Protein Pathways:

Proteasome

Product images:



Coomassie blue staining of purified PSMF1 protein (Cat# [TP315482]). The protein was produced from HEK293T cells transfected with PSMF1 cDNA clone (Cat# [RC215482]) using MegaTran 2.0 (Cat# [TT210002]).