

Product datasheet for PH317929

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PLAUR (NM 001005376) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: PLAUR MS Standard C13 and N15-labeled recombinant protein (NP 001005376)

Species: Human **HEK293 Expression Host: Expression cDNA Clone**

or AA Sequence:

RC217929

Predicted MW: 31.26 kDa

>RC217929 representing NM_001005376 **Protein Sequence:**

Red=Cloning site Green=Tags(s)

MGHPPLLPLLLLHTCVPASWGLRCMQCKTNGDCRVEECALGQDLCRTTIVRLWEEGEELELVEKSCTHS EKTNRTLSYRTGLKITSLTEVVCGLDLCNQGNSGRAVTYSRSRYLECISCGSSDMSCERGRHQSLQCRSP EEQCLDVVTHWIQEGEEGRPKDDRHLRGCGYLPGCPGSNGFHNNDTFHFLKCCNTTKCNEGPILELENLP QNGRQCYSCKGNSTHGCSSEETFLIDCRGPMNQCLVATGTHERSLWGSWLPCKSTTALRPPCCEEAQATH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

Labeling Method: Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

NP 001005376 RefSeq:

RefSeq Size: 1437 RefSeq ORF: 843

Synonyms: CD87; U-PAR; UPAR; URKR

Locus ID: 5329





UniProt ID: Q03405

Cytogenetics: 19q13.31

Summary: This gene encodes the receptor for urokinase plasminogen activator and, given its role in

localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the


extracellular matrix. It binds both the proprotein and mature forms of urokinase

plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage

reactions that have not yet been fully defined. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Secreted Protein **Protein Pathways:** Complement and coagulation cascades

Product images:



Coomassie blue staining of purified PLAUR protein (Cat# [TP317929]). The protein was produced from HEK293T cells transfected with PLAUR cDNA clone (Cat# [RC217929]) using MegaTran 2.0 (Cat# [TT210002]).