

Product datasheet for PH318795

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2D2 (NM 003339) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: UBE2D2 MS Standard C13 and N15-labeled recombinant protein (NP 003330)

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

or AA Sequence:

RC218795

Predicted MW: 16.7 kDa

>RC218795 protein sequence **Protein Sequence:**

Red=Cloning site Green=Tags(s)

MALKRIHKELNDLARDPPAQCSAGPVGDDMFHWQATIMGPNDSPYQGGVFFLTIHFPTDYPFKPPKVAFT TRIYHPNINSNGSICLDILRSQWSPALTISKVLLSICSLLCDPNPDDPLVPEIARIYKTDREKYNRIARE

WTQKYAM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

Labeling Method: Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

25 mM Tris-HCl, 100 mM glycine, pH 7.3 **Buffer:**

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 003330

RefSeg Size: 2715 RefSeq ORF: 441

Synonyms: E2(17)KB2; PUBC1; UBC4; UBC4/5; UBCH4; UBCH5B

Locus ID: 7322 UniProt ID: P62837





Cytogenetics: 5q31.2


Summary: Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs via

the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form polymeric chains, marking them as targets for 26S proteasome-mediated degradation. Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. This gene encodes a member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in multiple

transcript variants of this gene. [provided by RefSeq, May 2013]

Protein Pathways: Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified UBE2D2 protein (Cat# [TP318795]). The protein was produced from HEK293T cells transfected with UBE2D2 cDNA clone (Cat# [RC218795]) using MegaTran 2.0 (Cat# [TT210002]).