

# **Product datasheet for PH319382**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PKM2 (PKM) (NM 182470) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** PKM2 MS Standard C13 and N15-labeled recombinant protein (NP\_872270)

Species: Human **HEK293 Expression Host:** 

**Expression cDNA Clone** 

RC219382

or AA Sequence: Predicted MW:

**Protein Sequence:** 

57.9 kDa

>RC219382 representing NM\_182470

Red=Cloning site Green=Tags(s)

MSKPHSEAGTAFIQTQQLHAAMADTFLEHMCRLDIDSPPITARNTGIICTIGPASRSVETLKEMIKSGMN VARLNFSHGTHEYHAETIKNVRTATESFASDPILYRPVAVALDTKGPEIRTGLIKGSGTAEVELKKGATL KITLDNAYMEKCDENILWLDYKNICKVVEVGSKIYVDDGLISLQVKQKGADFLVTEVENGGSLGSKKGVN LPGAAVDLPAVSEKDIQDLKFGVEQDVDMVFASFIRKASDVHEVRKVLGEKGKNIKIISKIENHEGVRRF DEILEASDGIMVARGDLGIEIPAEKVFLAQKMMIGRCNRAGKPVICATQMLESMIKKPRPTRAEGSDVAN AVLDGADCIMLSGETAKGDYPLEAVRMQHLIAREAEAAMFHRKLFEELVRASSHSTDLMEAMAMGSVEAS YKCLAAALIVLTESGRSAHQVARYRPRAPIIAVTRNPQTARQAHLYRGIFPVLCKDPVQEAWAEDVDLRV

NFAMNVGKARGFFKKGDVVIVLTGWRPGSGFTNTMRVVPVP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

C-Myc/DDK Tag:

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Labeling Method:** Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stable for 3 months from receipt of products under proper storage and handling conditions. Stability:

RefSeq: NP 872270

RefSeq Size: 2674 RefSeq ORF: 1593





#### PKM2 (PKM) (NM\_182470) Human Mass Spec Standard - PH319382

Synonyms: CTHBP; HEL-S-30; OIP3; p58; PK3; PKM2; TCB; THBP1

**Locus ID:** 5315

**UniProt ID:** <u>P14618</u>, <u>A0A024R5Z9</u>

Cytogenetics: 15q23

**Summary:** This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase

that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP,

generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial

pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms


have been reported. [provided by RefSeq, May 2011]

**Protein Families:** Druggable Genome

**Protein Pathways:** Glycolysis / Gluconeogenesis, Metabolic pathways, Purine metabolism, Pyruvate metabolism,

Type II diabetes mellitus

## **Product images:**



Coomassie blue staining of purified PKM protein (Cat# [TP319382]). The protein was produced from HEK293T cells transfected with PKM cDNA clone (Cat# [RC219382]) using MegaTran 2.0 (Cat# [TT210002]).