

Product datasheet for PH320009

FGFR1 (NM_023106) Human Mass Spec Standard

Product data:

Product Type:	Mass Spec Standards
Description:	FGFR1 MS Standard C13 and N15-labeled recombinant protein (NP_075594)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC220009
Predicted MW:	81.9 kDa
Protein Sequence:	>RC220009 protein sequence Red=Cloning site Green=Tags(s)

MWSWKCLLFWAVLVATLCTARSPPTLPEQDALPSEDDDDDDSSSEKETDNTKPNPVAPYWT SPEKM
EKKLHAVPAAKTVKFKCPSGTPNPTRLRWLKNKKEFKPDHRIGGYKVRYATWSIIMDSVVPSPDKGNYTCI
VENEYGSINHTYQLDVVERSPHRPILQAGLPANKTVALGNSVEFMCKVYSDPQPHIQWLKHIEVNGSKIG
PDNLPPYQILKTAGVNTTDKEMEVLHLRNVSFEDAGEYTCLAGNSIGLSHHSAWLTVLEALEERPAMVTS
PLYLEIIYCTGAFLISCMVGSVIVYKMKSGTKKSDFFHSQMAVHKLAKSIPLRRQVTVSADSSASMNSGV
LLVRPSRLSSSGTPMLAGVSEYELPEDPRWELPRDRLVLGKPLGEGCFGQVVLAEIAGLDKDKPNRVTKV
AVKMLKSDATEKDLSDLISEMEMMKMIGKHKNIINLLGACTQDGPLYVIVEYASKGNLREYLQARRPPGL
EYCYNPSHNPEEQQLSSKDLVSCAYQVARGMEYLASKKCIHRDLAARNVLVTEEDNVMKIADFGIARDIHHI
DYKKTNGRLPVKWMPEALFDRIYTHQSDVWSFGVLLWEIFTLGGSPYPGVPVEELFKLLKEGHRMDK
PSNCTNELYMMRDCWHAVPSQRPTFKQLVEDLDRIVALTSNQEYLDLSMPLDQYSPSPDTRSTCSSG
EDSVFSHEPLPEEPCLPRHPAQLANGGLKRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	NP_075594



[View online »](#)


RefSeq Size:	5644
RefSeq ORF:	2193
Synonyms:	bFGF-R-1; BFGFR; CD331; CEK; ECCL; FGFR; FGFR-1; FLG; FLT-2; FLT2; HBGFR; HH2; HRTFDS; KAL2; N-SAM; OGD
Locus ID:	2260
UniProt ID:	P11362
Cytogenetics:	8p11.23

Summary: The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysplasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are associated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Adherens junction, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton

Product images:



Coomassie blue staining of purified FGFR1 protein (Cat# [TP320009]). The protein was produced from HEK293T cells transfected with FGFR1 cDNA clone (Cat# [RC220009]) using MegaTran 2.0 (Cat# [TT210002]).