

## **Product datasheet for PH320522**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PML Protein (PML) (NM 033246) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** PML MS Standard C13 and N15-labeled recombinant protein (NP\_150249)

Species: Human **HEK293 Expression Host:** 

**Expression cDNA Clone** 

RC220522

or AA Sequence: Predicted MW:

**Protein Sequence:** 

47.4 kDa

>RC220522 representing NM\_033246

Red=Cloning site Green=Tags(s)

MEPAPARSPRPQQDPARPQEPTMPPPETPSEGRQPSPSPSPTERAPASEEEFQFLRCQQCQAEAKCPKLL PCLHTLCSGCLEASGMQCPICQAPWPLGADTPALDNVFFESLQRRLSVYRQIVDAQAVCTRCKESADFWC FECEQLLCAKCFEAHQWFLKHEARPLAELRNQSVREFLDGTRKTNNIFCSNPNHRTPTLTSIYCRGCSKP LCCSCALLDSSHSELKCDISAEIQQRQEELDAMTQALQEQDSAFGAVHAQMHAAVGQLGRARAETEELIR ERVROVVAHVRAQERELLEAVDARYQRDYEEMASRLGRLDAVLQRIRTGSALVQRMKCYASDQEVLDMHG FLRQALCRLRQEEPQSLQAAVRTDGFDEFKVRLQDLSSCITQGKDAAVSKKASPEAASTPRDPIDVDLRN

**SGPTRTRRL**EQKLISEEDLAANDILDYKDDDDK**V** 

C-Myc/DDK Tag:

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Labeling Method:** Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stable for 3 months from receipt of products under proper storage and handling conditions. Stability:

RefSeq: NP 150249

RefSeg Size: 1851 RefSeq ORF: 1269

Synonyms: MYL; PP8675; RNF71; TRIM19





Locus ID: 5371

 UniProt ID:
 P29590

 Cytogenetics:
 15q24.1

Summary: The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM


motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to oncogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Extensive alternative splicing of this gene results in several variations of the protein's central and C-terminal regions; all variants encode the same N-terminus. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul

2008]

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Acute myeloid leukemia, Pathways in cancer, Ubiquitin mediated proteolysis

## **Product images:**



Coomassie blue staining of purified PML protein (Cat# [TP320522]). The protein was produced from HEK293T cells transfected with PML cDNA clone (Cat# [RC220522]) using MegaTran 2.0 (Cat# [TT210002]).