

Product datasheet for PH321453

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP6V1G3 (NM 133262) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: ATP6V1G3 MS Standard C13 and N15-labeled recombinant protein (NP_573569)

Species:HumanExpression Host:HEK293

Expression cDNA Clone

or AA Sequence:

RC221453

Predicted MW: 13.9 kDa

Protein Sequence: >RC221453 protein sequence

Red=Cloning site Green=Tags(s)

MTSQSQGIHQLLQAEKRAKDKLEEAKKRKGKRLKQAKEEAMVEIDQYRMQRDKEFRLKQSKIMGSQNNLS

DEIEEQTLGKIQELNGHYNKYMESVMNQLLSMVCDMKPEIHVNYRATN

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 573569

RefSeq Size: 645 RefSeq ORF: 354

Synonyms: ATP6G3; Vma10

 Locus ID:
 127124

 UniProt ID:
 Q96LB4

 Cytogenetics:
 1q31.3






Summary:

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'' and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes one of three G subunit proteins. Transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Pathways:

Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

Product images:



Coomassie blue staining of purified ATP6V1G3 protein (Cat# [TP321453]). The protein was produced from HEK293T cells transfected with ATP6V1G3 cDNA clone (Cat# [RC221453]) using MegaTran 2.0 (Cat# [TT210002]).