

Product datasheet for PH323161

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MATK (NM_139355) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: MATK MS Standard C13 and N15-labeled recombinant protein (NP_647612)

Species:HumanExpression Host:HEK293

Expression cDNA Clone

RC223161

or AA Sequence: Predicted MW:

56.3 kDa

Protein Sequence: >RC223161 representing NM_139355

Red=Cloning site Green=Tags(s)

MAGRGSLVSWRAFHGCDSAEELPRVSPRFLRAWHPPPVSARMPTRRWAPGTQCITKCEHTRPKPGELAFR KGDVVTILEACENKSWYRVKHHTSGQEGLLAAGALREREALSADPKLSLMPWFHGKISGQEAVQQLQPPE DGLFLVRESARHPGDYVLCVSFGRDVIHYRVLHRDGHLTIDEAVFFCNLMDMVEHYSKDKGAICTKLVRP KRKHGTKSAEEELARAGWLLNLQHLTLGAQIGEGEFGAVLQGEYLGQKVAVKNIKCDVTAQAFLDETAVM TKMQHENLVRLLGVILHQGLYIVMEHVSKGNLVNFLRTRGRALVNTAQLLQFSLHVAEGMEYLESKKLVH RDLAARNILVSEDLVAKVSDFGLAKAERKGLDSSRLPVKWTAPEALKHGKFTSKSDVWSFGVLLWEVFSY GRAPYPKMSLKEVSEAVEKGYRMEPPEGCPGPVHVLMSSCWEAEPARRPPFRKLAEKLARELRSAGAPAS

VSGQDADGSTSPRSQEP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 647612

RefSeq Size: 2183 RefSeq ORF: 1521





MATK (NM_139355) Human Mass Spec Standard - PH323161

Synonyms: CHK; CTK; HHYLTK; HYL; HYLTK; Lsk

Locus ID: 4145

UniProt ID: <u>P42679</u>, <u>F1T0G6</u>

Cytogenetics: 19p13.3


Summary: The protein encoded by this gene has amino acid sequence similarity to Csk tyrosine kinase

and has the structural features of the CSK subfamily: SRC homology SH2 and SH3 domains, a catalytic domain, a unique N terminus, lack of myristylation signals, lack of a negative regulatory phosphorylation site, and lack of an autophosphorylation site. This protein is thought to play a significant role in the signal transduction of hematopoietic cells. It is able to phosphorylate and inactivate Src family kinases, and may play an inhibitory role in the control of T-cell proliferation. This protein might be involved in signaling in some cases of breast cancer. Three alternatively spliced transcript variants that encode different isoforms have

been described for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Product images:



Coomassie blue staining of purified MATK protein (Cat# [TP323161]). The protein was produced from HEK293T cells transfected with MATK cDNA clone (Cat# [RC223161]) using MegaTran 2.0 (Cat# [TT210002]).