

Product datasheet for PH323668

CRYBA2 (NM_057093) Human Mass Spec Standard

Product data:

Product Type:	Mass Spec Standards
Description:	CRYBA2 MS Standard C13 and N15-labeled recombinant protein (NP_476434)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC223668
Predicted MW:	22.1 kDa
Protein Sequence:	>RC223668 protein sequence Red=Cloning site Green=Tags(s) MSSAPAPGPAPASLTLWDEEDFQGRRCRLLSDCANVCERGGLPRVRSVKVENVVWVAFEYPDFQGQQFIL EKGDYPRWSAWSGSSSHNSNQLLSFRPVLCANHNSRVTLFEGDNFQGCKFDLVDDYPSLPSMGWASKDV GSLKVSSGAWWAYQYPGYRGYQYVLERDRHSGEFCTYGELGTQAHTGQLQSIRRVQH TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	NP_476434
RefSeq Size:	903
RefSeq ORF:	591
Synonyms:	CTRCT42
Locus ID:	1412
UniProt ID:	P53672 , A0A024R429




[View online »](#)

Cytogenetics: 2q35

Summary: Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of the vertebrate eye, which function to maintain the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also defined as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Beta-crystallins, the most heterogeneous, differ by the presence of the C-terminal extension (present in the basic group but absent in the acidic group). Beta-crystallins form aggregates of different sizes and are able to form homodimers through self-association or heterodimers with other beta-crystallins. This gene is a beta acidic group member. Three alternatively spliced transcript variants encoding identical proteins have been reported. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified CRYBA2 protein (Cat# [TP323668]). The protein was produced from HEK293T cells transfected with CRYBA2 cDNA clone (Cat# [RC223668]) using MegaTran 2.0 (Cat# [TT210002]).