

## **Product datasheet for PH324067**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PIG3 (TP53I3) (NM\_147184) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** TP53I3 MS Standard C13 and N15-labeled recombinant protein (NP\_671713)

Species: Human
Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

RC224067

Predicted MW: 35.5 kDa

Protein Sequence: >RC224067 protein sequence

Red=Cloning site Green=Tags(s)

MLAVHFDKPGGPENLYVKEVAKPSPGEGEVLLKVAASALNRADLMQRQGQYDPPPGASNILGLEASGHVA ELGPGCQGHWKIGDTAMALLPGGGQAQYVTVPEGLLMPIPEGLTLTQAAAIPEAWLTAFQLLHLVGNVQA GDYVLIHAGLSGVGTAAIQLTRMAGAIPLVTAGSQKKLQMAEKLGAAAGFNYKKEDFSEATLKFTKGAGV NLILDCIGGSYWEKNVNCLALDGRWVLYGLMGGGDINGPLFSKLLFKRGSLITSLLRSRDNKYKQMLVNA

FTEQILPHFSTEGPQRLLPVLDRIYPVTEIQEAHKYMEANKNIGKIVLELPQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

**Storage:** Store at -80°C. Avoid repeated freeze-thaw cycles.

**Stability:** Stable for 3 months from receipt of products under proper storage and handling conditions.

**RefSeq:** NP 671713

RefSeq Size: 1643
RefSeq ORF: 996
Synonyms: PIG3
Locus ID: 9540





UniProt ID: Q53FA7

**Cytogenetics:** 2p23.3

**Summary:** The protein encoded by this gene is similar to oxidoreductases, which are enzymes involved

in cellular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppressor p53 and is thought to be involved in p53-mediated cell death. It contains a p53


consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucleotide microsatellite sequence. The microsatellite

is polymorphic, with a varying number of pentanucleotide repeats directly correlated with the extent of transcriptional activation by p53. It has been suggested that the microsatellite polymorphism may be associated with differential susceptibility to cancer. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, May 2011]

Protein Families: Druggable Genome
Protein Pathways: p53 signaling pathway

## **Product images:**



Coomassie blue staining of purified TP53I3 protein (Cat# [TP324067]). The protein was produced from HEK293T cells transfected with TP53I3 cDNA clone (Cat# [RC224067]) using MegaTran 2.0 (Cat# [TT210002]).