

Product datasheet for PH325257

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NACA (NM 001113202) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

NACA MS Standard C13 and N15-labeled recombinant protein (NP 001106673) **Description:**

Species: Human **HEK293 Expression Host: Expression cDNA Clone**

or AA Sequence:

RC225257

Predicted MW: 23.4 kDa

>Peptide sequence encoded by RC225257 **Protein Sequence:**

Blue=ORF Red=Cloning site Green=Tag(s)

MPGEATETVPATEQELPQPQAETGSGTESDSDESVPELEEQDSTQATTQQAQLAAAAEIDEEPVSKAKQ SRSEKKARKAMSKLGLRQVTGVTRVTIRKSKNILFVITKPDVYKSPASDTYIVFGEAKIEDLSQQAQLA AAEKFKVQGEAVSNIQENTQTPTVQEESEEEEVDETGVEVKDIELVMSQANVSRAKAVRALKNNSNDIV

NAIMELTM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Recombinant protein using RC225257 also available, TP325257M

C-Myc/DDK Tag:

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

Labeling Method: Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Store at -80°C. Avoid repeated freeze-thaw cycles. Storage:

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

NP 001106673 RefSeq:

RefSeq ORF: 645

Synonyms: HSD48; NAC-alpha; NACA1; skNAC

Locus ID: 4666

UniProt ID: Q13765, A0A024RB41






Cytogenetics: 12q13.3

Summary: This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form

the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum. This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes. [provided by RefSeq, Feb 2016]

Protein Families: Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified NACA protein (Cat# [TP325257]). The protein was produced from HEK293T cells transfected with NACA cDNA clone (Cat# [RC225257]) using MegaTran 2.0 (Cat# [TT210002]).