

# **Product datasheet for PH326567**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### NONO (NM\_001145408) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** NONO MS Standard C13 and N15-labeled recombinant protein (NP 001138880)

Species: Human
Expression Host: HEK293

Expression cDNA Clone

RC226567

or AA Sequence: Predicted MW:

54.2 kDa

Protein Sequence: >RC226567 protein sequence

Red=Cloning site Green=Tags(s)

MQSNKTFNLEKQNHTPRKHHQHHHQQQHHQQQQQPPPPPIPANGQQASSQNEGLTIDLKNFRKPGEKTF TQRSRLFVGNLPPDITEEEMRKLFEKYGKAGEVFIHKDKGFGFIRLETRTLAEIAKVELDNMPLRGKQLR VRFACHSASLTVRNLPQYVSNELLEEAFSVFGQVERAVVIVDDRGRPSGKGIVEFSGKPAARKALDRCSE GSFLLTTFPRPVTVEPMDQLDDEEGLPEKLVIKNQQFHKEREQPPRFAQPGSFEYEYAMRWKALIEMEKQ QQDQVDRNIKEAREKLEMEMEAARHEHQVMLMRQDLMRRQEELRRMEELHNQEVQKRKQLELRQEEERRR REEEMRRQQEEMMRRQQEGFKGTFPDAREQEIRMGQMAMGGAMGINNRGAMPPAPVPAGTPAPPGPATMM

PDGTLGLTPPTTERFGQAATMEGIGAIGGTPPAFNRAAPGAEFAPNKRRRY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 µg/µL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

**Storage:** Store at -80°C. Avoid repeated freeze-thaw cycles.

**Stability:** Stable for 3 months from receipt of products under proper storage and handling conditions.

**RefSeg:** NP 001138880

RefSeq Size: 3228 RefSeq ORF: 1413

Synonyms: MRXS34; NMT55; NRB54; P54; P54NRB; PPP1R114



#### NONO (NM\_001145408) Human Mass Spec Standard - PH326567

Locus ID: 4841

UniProt ID: <u>Q15233</u>, <u>A0A0S2Z4Z9</u>

Cytogenetics: Xq13.1


**Summary:** This gene encodes an RNA-binding protein which plays various roles in the nucleus, including

transcriptional regulation and RNA splicing. A rearrangement between this gene and the transcription factor E3 gene has been observed in papillary renal cell carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes exist on Chromosomes 2 and

16. [provided by RefSeq, Feb 2009]

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**



Coomassie blue staining of purified NONO protein (Cat# [TP326567]). The protein was produced from HEK293T cells transfected with NONO cDNA clone (Cat# [RC226567]) using MegaTran 2.0 (Cat# [TT210002]).