

Product datasheet for PH326678

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mortality Factor 4 like 2 (MORF4L2) (NM_001142421) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: MORF4L2 MS Standard C13 and N15-labeled recombinant protein (NP_001135893)

Species: Human Expression Host: HEK293

Expression cDNA Clone

RC226678

or AA Sequence: Predicted MW:

32.3 kDa

Protein Sequence: >RC226678 protein sequence

Red=Cloning site Green=Tags(s)

MSSRKQGSQPRGQQSAEEENFKKPTRSNMQRSKMRGASSGKKTAGPQQKNLEPALPGRWGGRSAENPPSG SVRKTRKNKQKTPGNGDGGSTSEAPQPPRKKRARADPTVESEEAFKNRMEVKVKIPEELKPWLVEDWDLV TRQKQLFQLPAKKNVDAILEEYANCKKSQGNVDNKEYAVNEVVAGIKEYFNVMLGTQLLYKFERPQYAEI LLAHPDAPMSQVYGAPHLLRLFVRIGAMLAYTPLDEKSLALLLGYLHDFLKYLAKNSASLFTASDYKVAS

AEYHRKAL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 001135893

RefSeq Size: 1929 RefSeq ORF: 864

Synonyms: MORFL2; MRGX

Locus ID: 9643





UniProt ID: <u>B3KP92</u>, <u>Q15014</u>

Cytogenetics: Xq22.2

Summary: Component of the NuA4 histone acetyltransferase complex which is involved in

transcriptional activation of select genes principally by acetylation of nucleosomal histone H4


and H2A. This modification may both alter nucleosome - DNA interactions and promote

interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the MSIN3A complex which acts to repress transcription by deacetylation of nucleosomal histones.

[UniProtKB/Swiss-Prot Function]

Protein Families: Transcription Factors

Product images:



Coomassie blue staining of purified MORF4L2 protein (Cat# [TP326678]). The protein was produced from HEK293T cells transfected with MORF4L2 cDNA clone (Cat# [RC226678]) using

MegaTran 2.0 (Cat# [TT210002]).