

Product datasheet for PP006B1

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

II1b Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB **Recommended Dilution:** ELISA:

To detect mIL-1 beta by direct ELISA (using 100 µl/well antibody solution) this antibody can be

used at a concentration of 0.15-0.30 µg/ml. Used in conjunction with compatible secondary

reagents, allows the detection of at least 0.2 ng/well of recombinant mIL-1 beta.

Western Blot:

To detect mIL-1 beta by Western Blot analysis this antibody can be used at a concentration of $0.1-0.2 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-1 beta is $1.5-3.0 \mu g/lane$, under either reducing or non-reducing conditions.

Reactivity: Mouse
Host: Rabbit
Clonality: Polyclonal

Immunogen: Highly pure (>98%) recombinant mIL-1 beta.

Specificity: Detect Murine IL-1 beta.

Formulation: PBS, pH 7.2 without preservatives.

Label: Biotin

State: Lyophilized purified Ig fraction.

Label: conjugated

Reconstitution Method: Restore in sterile PBS containing 0.1% BSA to a concentration

of 0.1-1.0 mg/ml.

Purification: Affinity chromatography.

Conjugation: Biotin

Storage: Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can

be stored at 2-8°C for one month or at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: interleukin 1 beta





Il1b Rabbit Polyclonal Antibody - PP006B1

Database Link: Entrez Gene 16176 Mouse

P10749

Background: Interleukin 1 (IL1), originally known as lymphocyte activating factor (LAF), activates T cells and

lymphocytes, which then proliferate and secrete interleukin 2. IL1 is primarily released from stimulated macrophages and monocytes, but also is released from several other cell types, and is thought to play a key role in inflammatory and immune responses. The two closely related agents, interleukin1 alpha (IL1 alpha) and interleukin1 beta (IL1 beta) bind to the same cell surface receptor, elicit nearly identical biological responses and share 25%

homology in their amino acid sequence.

Synonyms: IL-1 beta, IL1B, IL1 beta, IL1F2, Catabolin

Note: Centrifuge vial prior to opening!