

Product datasheet for PP015B2

Ccl7 Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Direct ELISA: To detect Mouse MCP-3 by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of ~1.0 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 2000-4000 pg/well of recombinant Mouse MCP-3. Sandwich ELISA: To detect Mouse MCP-3 by Sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This Biotin conjugated antibody, in conjunction with Purified Anti-Murine MCP-3 (Cat.-No PP015P) as a Capture antibody, allows the detection of at least 2000-4000 pg/well of recombinant mMCP-3. Western Blot: To detect mMCP-3 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mMCP-3 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) E.coli derived recombinant Murine MCP-3 (Cat.-No P024)
Specificity:	Reacts with Mouse Macrophage Chemotactic Protein-3 (MCP-3). Other species not tested.
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized, sterile filtered purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography.
Conjugation:	Biotin



[View online »](#)

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.
After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term.
Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: chemokine (C-C motif) ligand 7


Database Link: [Entrez Gene 20306 Mouse Q03366](#)

Background: This is a secreted chemokine which attracts macrophages during inflammation and metastasis. It is a member of the C-C subfamily of chemokines which are characterized by having two adjacent cysteine residues. The protein is an in vivo substrate of matrix metalloproteinase 2, an enzyme which degrades components of the extracellular matrix.

Synonyms: Small-inducible cytokine A7, CCL-7, MCP-3, C-C motif chemokine 7, SCYA6, SCYA7, NC28

Note: Centrifuge vial prior to opening!

Product images:



Sandwich ELISA using PP015B