

## Product datasheet for **PP033B1**

### Ifng Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	<u>ELISA:</u> To detect Rat IFN-gamma by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of 0.15-0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant Rat IFN-gamma. <u>Western Blot:</u> To detect Rat IFN-gamma by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IFN-gamma is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant rIFN gamma.
Specificity:	Recognises Rat IFN-gamma.
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	interferon gamma



[View online »](#)

**Database Link:** [Entrez Gene 25712 Rat P01581](#)

**Background:** Interferon gamma is mainly produced by T lymphocytes and NK cells. It is a pleiotropic cytokine involved in the regulation of nearly all phases of immune and inflammatory responses, including the activation, growth and differentiation of T cell, B cells, macrophages, NK cells and other cell types such as endothelial cells and fibroblasts. It has weak antiviral and antiproliferative activity, and potentiates the antiviral and anti tumor effects of IFN alpha / beta (type I interferon). It is upregulated by IL2, FGF basic, EGF and downregulated by vitamin D3 or DMN. Labile at pH 2.

**Synonyms:** IFN-gamma, gamma IFN

**Note:** Centrifuge vial prior to opening!