

Product datasheet for **PP1003B1**

CXCL5 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: (Direct): To detect hENA-78 by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant hENA-78. (Sandwich): To detect hENA-78 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human ENA-78 (PP1003P) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hENA-78. Western Blot: To detect hENA-78 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hENA-78 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hENA-78 (human ENA-78).
Specificity:	This antibody reacts with human Epithelial Neutrophil Activity Protein-78 (hENA-78)
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction.
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)

Stability:	Shelf life: One year from despatch.
Gene Name:	C-X-C motif chemokine ligand 5
Database Link:	Entrez Gene 6374 Human P42830
Background:	CXCL5 is a a 9.9kD inflammatory chemokine that belongs to the CXC chemokine family. CXCL5 is expressed by fibroblasts, is induced by bacterial lipopolysaccharides, and is chemotactic for neutrophils. This chemokine is produced concomitantly with Interleukin 8 (IL8) in response to stimulation with either Interleukin 1 (IL1) or Tumor Necrosis Factor alpha (TNFA). This chemokine is a potent chemotaxin involved in neutrophil activation.
Synonyms:	C-X-C motif chemokine 5, Small-inducible cytokine B5, ENA-78, SCYB5, Cytokine LIX
Note:	Centrifuge vial prior to opening!