

## Product datasheet for **PP1036B2**

### IL16 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: To detect hIL-16 by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of 0.15 - 0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant hIL-16. Western Blot: To detect hIL-16 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-16 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hIL-16 (human Interleukin 16).
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	interleukin 16
Database Link:	<a href="#">Entrez Gene 3603 Human Q14005</a>



[View online »](#)

**Background:**

IL16 is a proinflammatory cytokine that is chemotactic for CD4+ T lymphocytes, monocytes, and eosinophils. IL16 is synthesized as a precursor molecule (pro IL16) of approximately 68 kDa and 631 amino acid residues lacking a signal peptide. The sequence and structure of IL16 is conserved across species. In addition to inducing chemotaxis, IL16 upregulates the IL2 receptor and upregulates HLADR expression. It also inhibits T cell receptor (TCR)/CD3 dependent activation, and suppresses HIV1 replication in vitro. Sources of IL16 include epithelial cells, mast cells, T lymphocytes (CD4+ and CD8+), macrophages, synovial fibroblasts, and eosinophils.

**Synonyms:**

IL-16, LCF

**Note:**

Centrifuge vial prior to opening!