

Product datasheet for **PP1039B1**

IP10 (CXCL10) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: To detect hIP-10 by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of 0.15 - 0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant hIP-10. Western Blot: To detect hIP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIP-10 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hIP-10 (human IP-10).
Specificity:	Human Interferon-gamma Inducible Protein-10 (IP-10).
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	C-X-C motif chemokine ligand 10
Database Link:	Entrez Gene 3627 Human P02778



[View online »](#)

- Background:** Interferon-gamma-inducible 10 kD protein (IP-10), is a CXC chemokine with chemoattractant properties for CD4-positive T cells and inhibits early normal and leukemic hemopoietic progenitor proliferation. IP-10 is produced by a wide variety of cell types ranging from neutrophils and monocytes to hepatocytes, endothelial cells and keratinocytes. The cytokine is reported to be involved in a scala of inflammatory pathologies such as HIV encephalitis, cutaneous T cell lymphoma, chronic hepatitis and acute anterior uveitis. Various observations strongly suggest a role for the CXC chemokines IL-8 and IP-10 in the regulation of angiogenic activity in cancer and in idiopathic pulmonary fibrosis.
- Synonyms:** C-X-C motif chemokine 10, Small-inducible cytokine B10, Gamma-IP10, INP10, SCYB10
- Note:** Centrifuge vial prior to opening!