

Product datasheet for **PP1054B2**

Macrophage Inflammatory Protein 4 (CCL18) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: To detect hMIP-4 by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of 0.15 - 0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant hMIP-4 Western Blot: To detect hMIP-4 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIP-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hMIP-4.
Specificity:	Specific for human Macrophage Inflammatory Protein-4 (hMIP-4) (PARC).
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	C-C motif chemokine ligand 18
Database Link:	Entrez Gene 6362 Human P55774



[View online »](#)

Background:

This chemotactic factor that attracts lymphocytes but not monocytes or granulocytes may be involved in B cell migration into B cell follicles in lymph nodes. It attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes, has chemotactic activity for naive T cells, CD4+ and CD8+ T cells and thus may play a role in both humoral and cell-mediated immunity responses. It is expressed at high levels in lung, lymph nodes, placenta, and bone marrow, not expressed by peripheral blood monocytes, and a monocyte-to-macrophage differentiation is a prerequisite for expression. Specifically induced in macrophages by IL4, IL13, and IL10, it belongs to the intercrine beta (chemokine CC) family.

Synonyms:

C-C motif chemokine 18, AMAC1, DCCK1, MIP-4, PARC, Small-inducible cytokine A18, SCYA18

Note:

Centrifuge vial prior to opening!