

Product datasheet for PP1056B2

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CXCL7 (PPBP) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: ELISA: To detect hNAP-2 by direct ELISA (using 100 μl/well antibody solution) this antibody can

be used at a concentration of 0.15 - 0.30 μ g/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant hNAP-2. Western Blot: To detect hNAP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNAP-2 is 1.5 - 3.0 ng/lane, under either reducing or non-

reducing conditions.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Highly pure (>98%) recombinant hNAP-2.

Specificity: Human Neutrophil Activating Protein-2 (hNAP-2).

Formulation: PBS, pH 7.2 without preservatives.

Label: Biotin

State: Lyophilized purified Ig fraction.

Label: conjugated

Reconstitution Method: Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Purification: Affinity chromatography.

Conjugation: Biotin

Storage: Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can

be stored at 2-8°C for one month or at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: pro-platelet basic protein

Database Link: Entrez Gene 5473 Human

P02775





CXCL7 (PPBP) Rabbit Polyclonal Antibody - PP1056B2

Background: NAP2 is a platelet-derived growth factor that belongs to the CXC chemokine family. This

growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and sythesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator

by synovial cells.

Synonyms: CTAP3, SCYB7, TGB1, THBGB1

Note: Centrifuge vial prior to opening!