

Product datasheet for **PP1065B2**

Neurturin (NRTN) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: Direct: To detect hNeurturin (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This In conjunction with compatible secondary reagents, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hNeurturin. Sandwich: To detect hNeurturin (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. In conjunction with Polyclonal Anti-Human Neurturin as a capture antibody, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hNeurturin. Western blot: To detect hNeurturin this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNeurturin is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (> 98 %) recombinant human Neurturin
Specificity:	This antibody detects Neurturin.
Formulation:	PBS, pH 7.2 Label: Biotin State: Sterile filtered lyophilized Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile PBS containing 0.1 % BSA to a concentration of 0.1 - 1.0 mg/ml.
Purification:	Affinity chromatography
Conjugation:	Biotin
Storage:	Store the lyophilized antibody at -20 °C. Following reconstitution it is stable for two weeks at 2 - 8 °C. Frozen aliquots are stable for 6 months when stored at -20 °C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.



[View online »](#)

Gene Name: neurturin

Database Link: [Entrez Gene 4902 Human Q99748](#)

Background: Human Neurturin (NTN) and Human GDNF comprise a family of TGF-beta related neurotrophic factor which have trophic influences on a variety of neuronal populations. Neurturin promotes the survival of certain sympathetic and sensory neurins through interaction with distinct set of GDNF-like receptors.

Synonyms: NRTN