

## Product datasheet for **PP1069B1**

### TARC (CCL17) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<b>ELISA:</b> <u>Direct:</u> To detect hTARC by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant hTARC. <u>Sandwich:</u> To detect hTARC by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human TARC (PP1069P) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hTARC. <b>Western Blot:</b> To detect hTARC by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTARC is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hTARC.
Specificity:	This antibody reacts with TARC (CCL17).
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized (sterile filtered) purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Biotin



[View online »](#)

<b>Storage:</b>	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: One year from despatch.
<b>Gene Name:</b>	C-C motif chemokine ligand 17
<b>Database Link:</b>	<a href="#">Entrez Gene 6361 Human Q92583</a>
<b>Synonyms:</b>	C-C motif chemokine 17, Small-inducible cytokine A17, SCYA17
<b>Note:</b>	Centrifuge vial prior to opening!