

# **Product datasheet for PP1074P1**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Thrombopoietin (THPO) Goat Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ELISA, FN, WB

**Recommended Dilution:** Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hTPO

(5.00 ng/ml), a concentration of 4.2-6.3 μg/ml of this antibody is required.

ELISA:

(Indirect): To detect httpO by indirect ELISA (using 100 µl/well antibody solution) a concentration of 0.5-2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least

0.2-0.4 ng/well of recombinant hTPO.

(Sandwich): To detect hTPO by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5-2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human TPO (PP1074B) as a detection

antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hTPO. Western Blot: To detect hTPO by Western Blot analysis this antibody can be used at a

concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTPO is 1.5-3.0 ng/lane, under either reducing or non-

reducing conditions.

Reactivity: Human Host: Goat

Clonality: Polyclonal

Immunogen: Highly purified recombinant human TPO.Specificity: This antibody reacts with Thrombopoietin.

**Formulation:** PBS, pH 7.2 without preservatives.

State: Aff - Purified

State: Lyophilized purified Ig fraction.

**Reconstitution Method:** Restore in sterile water to a concentration of 0.1-1.0 mg/ml.

**Purification:** Affinity chromatography.

Conjugation: Unconjugated





### Thrombopoietin (THPO) Goat Polyclonal Antibody - PP1074P1

**Storage:** Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can

be stored at 2-8°C for one month or at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch.

Gene Name: thrombopoietin

Database Link: Entrez Gene 7066 Human

P40225

Synonyms: THPO, MGDF, C-mpl ligand

**Note:** Centrifuge vial prior to opening!