

Product datasheet for **PP1083P2**

Cxcl10 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, FN, WB
Recommended Dilution:	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Mouse IP-10 (100 ng/ml), a concentration of 10.0 µg/ml of this antibody is required. Sandwich ELISA: To detect Mouse IP-10 by Sandwich ELISA (using 100 µl/well antibody solution), a concentration of 0.5-2.0 µg/ml of this antibody is required. This Affinity purified antibody, in conjunction with Biotin conjugated anti Mouse IP10 antibody (cat. PP1083B) as a Detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant Mouse IP-10. Western Blot: To detect Mouse IP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Mouse IP-10 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) E.coli derived recombinant Murine IP-10.
Specificity:	Reacts with Interferon-gamma Inducible Protein.
Formulation:	PBS, pH 7.2 without preservatives. State: Aff - Purified State: Lyophilized purified Ig fraction.
Reconstitution Method:	Restore in sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography using an immobilized Murine IP-10 Matrix
Conjugation:	Unconjugated
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	chemokine (C-X-C motif) ligand 10



[View online »](#)


Database Link: [Entrez Gene 15945 Mouse P17515](#)

Background: Interferon-gamma-inducible 10 kD protein (IP-10), is a CXC chemokine with chemoattractant properties for CD4-positive T cells and inhibits early normal and leukemic hemopoietic progenitor proliferation. IP-10 is produced by a wide variety of cell types ranging from neutrophils and monocytes to hepatocytes, endothelial cells and keratinocytes. The cytokine is reported to be involved in a scala of inflammatory pathologies such as HIV encephalitis, cutaneous T cell lymphoma, chronic hepatitis and acute anterior uveitis. Various observations strongly suggest a role for the CXC chemokines IL-8 and IP-10 in the regulation of angiogenic activity in cancer and in idiopathic pulmonary fibrosis.

Synonyms: C-X-C motif chemokine 10, Small-inducible cytokine B10, Gamma-IP10, INP10, SCYB10

Note: Centrifuge vial prior to opening!

Product images:



To detect mIP-10 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with the Biotinylated Anti-Murine IP-10 [PP1083B1]/PP1083P2 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mIP-10.