

Product datasheet for **PP1116B2**

BCA1 (CXCL13) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<p>Direct ELISA: To detect hBCA-1 by direct ELISA (using 100 µl/well antibody solution) a concentration of ~1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hBCA-1.</p> <p>Sandwich ELISA: To detect hBCA-1 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human BCA-1 as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hBCA-1.</p> <p>Western Blot: To detect hBCA-1 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBCA-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.</p>
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) E.coli derived recombinant Human CXCL13 / BCA1
Specificity:	This antibody detects Human CXCL13 / BCA1.
Formulation:	<p>PBS, pH 7.2 without preservatives</p> <p>Label: Biotin</p> <p>State: Lyophilized (sterile filtered) purified Ig fraction</p>
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography
Conjugation:	Biotin
Storage:	<p>Prior to reconstitution store at 2-8°C.</p> <p>Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.</p> <p>Avoid repeated freezing and thawing.</p>



[View online »](#)

Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	150 kDa
Gene Name:	C-X-C motif chemokine ligand 13
Database Link:	Entrez Gene 10563 Human O43927
Background:	B lymphocyte chemoattractant, independently cloned and named Angie, is a CXC chemokine strongly expressed in the follicles of the spleen, lymph nodes, and Peyer's patches. It preferentially promotes the migration of B lymphocytes (compared to T cells and macrophages), apparently by stimulating calcium influx into, and chemotaxis of, cells expressing Burkitt's lymphoma receptor 1 (BLR1). It may therefore function in the homing of B lymphocytes to follicles.
Synonyms:	BLC, SCYB13, C-X-C motif chemokine 13
Note:	Centrifuge vial prior to opening!