

## Product datasheet for **PP1126B1**

### Egf Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA
Recommended Dilution:	ELISA: (Direct): To detect mEGF by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant mEGF. (Sandwich): To detect mEGF by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Murine EGF (PP1126P) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant mEGF. Western Blot: To detect mEGF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mEGF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant mEGF.
Specificity:	Murine EGF
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity chromatography
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)

<b>Stability:</b>	Shelf life: One year from despatch.
<b>Gene Name:</b>	epidermal growth factor
<b>Database Link:</b>	<a href="#">Entrez Gene 13645 Mouse P01132</a>
<b>Background:</b>	<p>Epidermal growth factor (EGF) has a profound effect on the differentiation of specific cells in vivo and is a potent mitogenic factor for a variety of cultured cells. The EGF precursor is believed to exist as a membrane bound molecule which is proteolytically cleaved to generate the 53 amino acid peptide hormone that stimulates cells to divide. EGF exerts its actions by binding to the EGFR, a 170 kDa protein.</p> <p>Epidermal growth factor (EGF) is a key growth factor regulating cell survival. Through its binding to cell surface receptors, EGF activates an extensive network of signal transduction pathways that include activation of the PI3K/AKT, RAS/ERK and JAK/STAT pathways. Because of its key role in driving the proliferation of cells, EGFR is a target of several anti-cancer drugs currently in development.</p>
<b>Synonyms:</b>	Urogastrone, Epidermal growth factor, URG, HOMG4
<b>Note:</b>	Centrifuge vial prior to opening!