

## Product datasheet for **PP1205B1**

### Cxcl16 Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: To detect mCXCL16 by direct ELISA (using 100 µl/well antibody solution) this antibody can be used at a concentration of 0.15-0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant mCXCL16. Western Blot: To detect mCXCL16 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mCXCL16 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant mCXCL16.
Formulation:	PBS, pH 7.4 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction.
Reconstitution Method:	Restore to a concentration of 50 µg/ml with sterile PBS solution containing 0.1% BSA
Purification:	Affinity chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	chemokine (C-X-C motif) ligand 16
Database Link:	<a href="#">Entrez Gene 66102 Mouse</a> <a href="#">Q8BSU2</a>



[View online »](#)

**Background:** CXCL16 acts as a scavenger receptor on macrophages, which specifically binds to OxLDL (oxidized low density lipoprotein), suggesting that it may be involved in pathophysiology such as atherogenesis. It induces a strong chemotactic response and calcium mobilization. It binds to CXCR6/Bonzo.

**Synonyms:** C-X-C motif chemokine 16, Small-inducible cytokine B16, SR-PSOX, SCYB16, SRPSOX