

## Product datasheet for **PP1206P2**

### Prokineticin 1 (PROK1) Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Neutralization. ELISA: To detect hEG-VEGF by direct ELISA (using 100 µl/well antibody solution) a concentration of at least 0.5 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2-0.4 ng/well of recombinant hEG-VEGF. Western Blot: To detect hEG-VEGF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEG-VEGF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Highly pure (>98%) recombinant hEG-VEGF.
Formulation:	0.5 X PBS pH 7.4 without preservatives. State: Aff - Purified State: Lyophilized purified Ig fraction.
Reconstitution Method:	Restore in sterile water to a concentration of > 0.2 mg/ml.
Purification:	Affinity chromatography.
Conjugation:	Unconjugated
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	prokineticin 1
Database Link:	<a href="#">Entrez Gene 84432 Human P58294</a>
Synonyms:	Prokineticin-1, Prokineticin 1, Mambakine, EG-VEGF



[View online »](#)