

## **Product datasheet for PP1213B2**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Midkine (MDK) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: ELISA: To detect hMidkine by direct ELISA (using 100 µl/well antibody solution) this antibody

can be used at a concentration of 0.15 - 0.30 µg/ml. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant hMidkine. Western Blot: To detect hMidkine by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMidkine is 1.5 - 3.0 ng/lane, under either reducing or non-

reducing conditions.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

**Immunogen:** Highly pure (>98%) recombinant hMidkine.

**Formulation:** PBS, pH 7.2 without preservatives.

Label: Biotin

State: Lyophilized purified Ig fraction.

Label: conjugated

**Reconstitution Method:** Restore to a concentration of 50 μg/ml with sterile PBS solution containing 0.1% BSA.

**Purification:** Affinity chromatography.

**Conjugation:** Biotin

**Storage:** Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can

be stored at 2-8°C for one month or at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** midkine (neurite growth-promoting factor 2)

**Database Link:** Entrez Gene 4192 Human

P21741





## Midkine (MDK) Rabbit Polyclonal Antibody - PP1213B2

**Background:** Midkine exhibits neurite outgrowth-promoting activity and may play a role in nervous system

development and/or maintenance. Its expression is predominant only for a short period from approximately one-half to two-thirds of the way through gestation; before and after that, it is barely detectable. Midkine was first found in differentiating mouse teratocarcinoma cells. It has neurotrophic activities and is mitogenic to certain, but not to all, fibroblast cell lines.

Synonyms: MK, MDK, MK1, NEGF2, Neurite outgrowth-promoting protein