

Product datasheet for PP1213P1

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Midkine (MDK) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: Neutralization.

ELISA: To detect hMidkine by direct ELISA (using 100 μl/well antibody solution) a

concentration of at least 0.5 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4

ng/well of recombinant hMidkine.

Western Blot: To detect hMidkine by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMidkine is 1.5 - 3.0 ng/lane, under either reducing or non-

reducing conditions.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Highly pure (>98%) recombinant hMidkine

Formulation: 0.5 X PBS pH 7.4 without preservatives.

State: Aff - Purified

State: Lyophilized purified Ig fraction.

Reconstitution Method: Restore in sterile water to a concentration of > 0.2 mg/ml.

Purification: Affinity chromatography employing immobilized hMidkine matrix.

Conjugation: Unconjugated

Storage: Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can

be stored at 2-8°C for one month or at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: midkine (neurite growth-promoting factor 2)

Database Link: Entrez Gene 4192 Human

P21741





Midkine (MDK) Rabbit Polyclonal Antibody - PP1213P1

Background: Midkine exhibits neurite outgrowth-promoting activity and may play a role in nervous system

development and/or maintenance. Its expression is predominant only for a short period from approximately one-half to two-thirds of the way through gestation; before and after that, it is barely detectable. Midkine was first found in differentiating mouse teratocarcinoma cells. It has neurotrophic activities and is mitogenic to certain, but not to all, fibroblast cell lines.

Synonyms: MK, MDK, MK1, NEGF2, Neurite outgrowth-promoting protein